MAXIMUS Federal Services

Case Number:	CM15-0113251		
Date Assigned:	06/19/2015	Date of Injury:	07/09/2008
Decision Date:	07/20/2015	UR Denial Date:	05/15/2015
Priority:	Standard	Application Received:	06/11/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: North Carolina
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 40-year-old male, who sustained an industrial injury on 7/9/08. The injured worker was diagnosed as having rule out lumbar spine disc displacement, lumbosacral neuritis/radiculitis and lumbar spine muscle spasm. Treatment to date has included lumbar fusion in 2011, physical therapy, oral medications including Norco, Soma, Gabapentin and Percocet; activity restrictions and home exercise program. Currently, the injured worker complains of continued back pain rated 10/10 without medications and 8/10 with medications and right leg numbness. Physical exam noted a limp; torso inclined forward, painful range of motion, weakness of right leg and tenderness of lumbar spine with spasm. A request for authorization was submitted for Norco, Percocet, Gabapentin, Soma and pain management.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

1 prescription of Norco 10/325mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain dairy that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work; (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant improvement in VAS scores for significant periods of time. There are no objective measurements of improvement in function. Therefore, not all criteria for the ongoing use of opioids have been met and the request is not medically necessary.

1 prescription of Soma 350mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines muscle relaxants Page(s): 63-65.

Decision rationale: The California chronic pain medical treatment guidelines section on muscle relaxants states: Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. In addition, there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) (Chou, 2004) This medication is not intended for long-term use per the California MTUS. The medication has not been prescribed for the flare-up of chronic low back pain. This is not an approved use for the medication. For these reasons, criteria for the use of this medication have not been met. Therefore, the request is not medically necessary.

1 Pain Management Consultation: Overturned

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Chronic Pain Disorder Medical Treatment Guidelines, State of Colorado Department of Labor and Employment (Chapter: Chronic Pain Disorder; Section: Therapeutic procedures. Non-Operative) 4/27/2007 pg 56.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 2 General Approach to Initial Assessment and Documentation, Chapter 3 Initial Approaches to Treatment.

Decision rationale: Per the ACOEM: The health practitioner may refer to other specialist if a diagnosis is uncertain or extremely complex, when psychosocial factors are present, or when the plan or course of care may benefit from additional expertise. A referral may be for 1. Consultation to aid in the diagnosis, prognosis, therapeutic management, determination of medical stability. The patient has ongoing complaints of ongoing pain that have failed treatment by the primary treating physician. Therefore, criteria for a pain management consult have been met and the request is medically necessary.