

Case Number:	CM15-0113633		
Date Assigned:	06/19/2015	Date of Injury:	08/14/2013
Decision Date:	07/21/2015	UR Denial Date:	05/14/2015
Priority:	Standard	Application Received:	06/11/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California
 Certification(s)/Specialty: Physical Medicine & Rehabilitation

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 55 year old male, who sustained an industrial injury on 8/14/2013. He reported falling backwards and experiencing low back pain. Diagnoses have included sciatica, displaced lumbar intervertebral disc, insomnia, anxiety and depression. Treatment to date has included trigger point injections, chiropractic treatment, magnetic resonance imaging (MRI) and medication. According to the progress report dated 5/4/2015, the injured worker complained of constant low back pain that traveled to the right leg and ankle. The pain was rated 9/10. Objective findings revealed weakness in the right leg. The injured worker could not toe or heel walk due to pain/weakness. He used a cane to ambulate. Authorization was requested for Norco, Soma and Alprazolam.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

Decision rationale: Per the MTUS Guidelines, cite opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury without acute flare, new injury, or progressive deterioration. The Norco 10/325mg #90 is not medically necessary and appropriate.

Soma 350mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Carisoprodol (Soma), page 29.

Decision rationale: Per MTUS Chronic Pain Guidelines on muscle relaxant, Soma is not recommended for mild to moderate chronic persistent pain problems including chronic pain (other than for acute exacerbations) due to the high prevalence of adverse effects in the context of insufficient evidence of benefit as compared to other medications. Guidelines do not recommend long-term use of this muscle relaxant for this chronic injury. Additionally, the efficacy in clinical trials has been inconsistent and most studies are small and of short duration. These medications may be useful for chronic musculoskeletal pain, but there are no long-term studies of their effectiveness or safety. Submitted reports have not adequately demonstrated the indication or medical need for this treatment and there is no report of progressive deterioration in clinical findings, acute flare-up or new injury to support for its long-term use. There is no report of functional improvement resulting from its previous treatment to support further use as the patient remains unchanged. The Soma 350mg #90 is not medically necessary and appropriate.

Alprazolam .5mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines, page 24.

Decision rationale: Alprazolam is indicated for the management of anxiety disorder. Anxiety or tension associated with the stress of everyday life usually does not require treatment with an anxiolytic. Alprazolam is an anti-anxiety medication in the benzodiazepine family which inhibits many of the activities of the brain as it is believed that excessive activity in the brain may lead to anxiety or other psychiatric disorders. Per the Chronic Pain Treatment Guidelines, benzodiazepines are not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks as chronic benzodiazepines are the treatment of choice in very few conditions and tolerance to hypnotic effects develops rapidly. Additionally, submitted reports have not demonstrated clear functional benefit of treatment already rendered. The Alprazolam .5mg #30 is not medically necessary and appropriate.