

Case Number:	CM15-0114250		
Date Assigned:	06/25/2015	Date of Injury:	07/12/2014
Decision Date:	07/24/2015	UR Denial Date:	06/05/2015
Priority:	Standard	Application Received:	06/15/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 60 year old man sustained an industrial injury on 7/12/2014 after being involved in a motor vehicle accident. Diagnoses include neck sprain and probable facet syndrome. Treatment has included oral medications and chiropractic care. Physician notes on a PR-2 dated 2/4/2015 show complaints of neck pain and intermittent headaches. Recommendations include continue chiropractic care and follow up in one month.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Anapox DS Naproxen sodium 550mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants Page(s): 63.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NON SELECTIVE NSAIDS Page(s): 72.

Decision rationale: Naproxen (Naprosyn): delayed release (EC-Naprosyn), as Sodium salt (Anaprox, Anaprox DS, Aleve [otc]) Generic available; extended-release (Napreelan): 375 mg.

Different dose strengths and formulations of the drug are not necessarily bioequivalent. Dosing Information: Osteoarthritis or ankylosing spondylitis: Dividing the daily dose into 3 doses versus 2 doses for immediate-release and delayed-release formulations generally does not affect response. Morning and evening doses do not have to be equal in size. The dose may be increased to 1500 mg/day of naproxyn for limited periods when a higher level of analgesic/anti-inflammatory activity is required (for up to 6 months). Naprosyn or naproxyn: 250-500 mg PO twice daily. Anaprox: 275-550 mg PO twice daily. (total dose may be increased to 1650 mg a day for limited periods). EC-Naprosyn: 375 mg or 500 mg twice daily. The tablet should not be broken, crushed or chewed to maintain integrity of the enteric coating. Naprelan: Two 375 mg tablets (750 mg) PO once daily or two 500 mg tablets (1000 mg) once daily. If required (and a lower dose was tolerated) Naprelan can be increased to 1500 mg once daily for limited periods (when higher analgesia is required). Pain: Naprosyn or naproxyn: 250-500 mg PO twice daily. The maximum dose on day one should not exceed 1250 mg and 1000 mg on subsequent days. Anaprox: 275-550 mg PO twice daily. The maximum dose on day one should not exceed 1375 mg and 1100 mg on subsequent days. Extended-release Naprelan: Not recommended due to delay in absorption. (Naprelan Package Insert) There is no documentation of the rationale behind the long-term use of Naproxen. NSAID should be used for the shortest duration and the lowest dose. There is no documentation from the patient's file that the provider titrated Naproxen to the lowest effective dose and used it for the shortest period possible. Naproxen was used without clear documentation of its efficacy. Furthermore, there is no documentation that the provider followed the patient for NSAID adverse reactions that are not limited to GI side effect, but also may affect the renal function. Therefore, the request for NAPROXEN 550 mg #90 is not medically necessary.

Fexmid Cyclobenzaprine 7.5mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants Page(s): 64.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 63.

Decision rationale: According to MTUS guidelines, Cyclobenzaprine a non sedating muscle relaxants is recommended with caution as a second line option for short term treatment of acute exacerbations in patients with chronic spasm and pain. Efficacy appears to diminish over time and prolonged use may cause dependence. The guidelines do not recommend to be used for more than 2-3 weeks. The patient in this case does not have clear recent evidence of spasm and the prolonged use of Cyclobenzaprine is not justified. Therefore, the request for Cyclobenzaprine 7.5mg #60 is not medically necessary.

(DOS 5/20/15) Ultram Tramadol HCL ER 150mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 39-94, 76-78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines
Tramadol Page(s): 113.

Decision rationale: According to MTUS guidelines, Ultram (Tramadol) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. Although, Tramadol may be needed to help with the patient pain, there is no clear evidence of objective and recent functional and pain improvement from its previous use. There is no objective documentation of pain severity level to justify the use of tramadol in this patient. There is no clear documentation of the efficacy/safety of previous use of tramadol. There is no recent evidence of objective monitoring of compliance of the patient with his medications. Therefore, the prescription of TRAMADOL HCL ER 150 mg #60 is not medically necessary.