

|                       |              |                              |            |
|-----------------------|--------------|------------------------------|------------|
| <b>Case Number:</b>   | CM15-0114709 |                              |            |
| <b>Date Assigned:</b> | 06/23/2015   | <b>Date of Injury:</b>       | 08/27/2014 |
| <b>Decision Date:</b> | 07/22/2015   | <b>UR Denial Date:</b>       | 05/27/2015 |
| <b>Priority:</b>      | Standard     | <b>Application Received:</b> | 06/15/2015 |

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: North Carolina  
 Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 50 year old male, who sustained an industrial injury on 8/27/14. The injured worker has complaints of neck pain. The documentation noted on 5/14/15 that the injured worker reported that he tore his left Achilles tendon and had minor tear. The documentation that he was running across the street and stopped suddenly and had excruciating pain, seen in the emergency room and was issued oxycodone 20 tabs for acute pain. The documentation noted that he has two pills left, was placed in a walking boot that he stopped two weeks prior and the pain is very minor now. The documentation noted that he was taking the medications or severe left ankle pain and notices that it helped to decrease his neck pain. The injured worker continues with neck pain radiating down to left hand and pinky and ring fingers with numbness/tingling and feels he is losing strength in fingers. The cervical spine range of motion is restricted with flexion limited to 30 degrees, extension limited to 30 degrees, right lateral bending limited to 37 degrees and left lateral bending limited to 35 degrees. The diagnoses have included cervical radiculopathy; cervical facet syndrome and cervical pain. Treatment to date has included lyrica; nucynta; ibuprofen; injections; home exercise program; transcutaneous electrical nerve stimulation unit; electromyography/nerve conduction velocity study of the left upper extremity and completed cervical magnetic resonance imaging (MRI) showed C5, C6 and C7 mild disc degeneration on 4/2/15. The request was for narc-oxycodone HCL 5 mg #60.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Narc-Oxycodone HCL 5 mg #60:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Specific Drug List.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines opioids  
Page(s): 76-84.

**Decision rationale:** The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work, (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox- AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant improvement in VAS scores for significant periods of time. There are no objective measurements of improvement in function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

