

Case Number:	CM15-0115416		
Date Assigned:	06/23/2015	Date of Injury:	01/30/2003
Decision Date:	10/05/2015	UR Denial Date:	05/29/2015
Priority:	Standard	Application Received:	06/15/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, District of Columbia, Maryland
 Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 63-year-old female, who sustained an industrial injury on 1-30-03. The injured worker has complaints of bilateral wrist, hand and forearm pain and neck, shoulder and trapezius pain, more on the right and cervical spine discomfort. The documentation noted slight tenderness over the distal forearm and volar wrist bilaterally, right greater than left. Shoulder examination revealed slight tenderness of the trapezius and right shoulder over the acromioclavicular region to a slight degree and left side is mildly tender. Cervical spine examination reveals slight tenderness and spasm noted mostly in the right paracervical region and extending into the right trapezius and right scapular region. The diagnoses have included overuse syndrome of both upper extremities with bilateral wrist, hand and forearm tendinitis, right greater than left; bilateral shoulder and trapezius strain, right worse than left and secondary cervical strain due to overuse, right worse than left. Treatment to date has included transcutaneous electrical nerve stimulation unit; home exercise program; therapy; ambien; norco; flexeril ; naproxen and ranitidine. The request was for norco 5-325mg #15; flexeril 5mg #30 ambien 10mg.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 5/325mg #15: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 78, 91.

Decision rationale: Per MTUS Chronic Pain Medical Treatment Guidelines p78 regarding ongoing management of opioids "Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: Pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug related behaviors. These domains have been summarized as the "4A's" (Analgesia, activities of daily living, adverse side effects, and any aberrant drug-taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs." Review of the available medical records reveals no documentation to support the medical necessity of norco nor any documentation addressing the '4A's' domains, which is a recommended practice for the on-going management of opioids. Specifically, the notes do not appropriately review and document pain relief, functional status improvement, appropriate medication use, or side effects. The MTUS considers this list of criteria for initiation and continuation of opioids in the context of efficacy required to substantiate medical necessity, and they do not appear to have been addressed by the treating physician in the documentation available for review. Furthermore, efforts to rule out aberrant behavior (e.g. CURES report, UDS, opiate agreement) are necessary to assure safe usage and establish medical necessity. There is no documentation comprehensively addressing this concern in the records available for my review. As MTUS recommends discontinuing opioids if there is no overall improvement in function, medical necessity cannot be affirmed.

Flexeril 5mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants Page(s): 63-64.

Decision rationale: With regard to muscle relaxants, the MTUS CPMTG states: "Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (Van Tulder, 2003) (Van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement." Regarding Cyclobenzaprine: "Recommended for a short course of therapy. Limited, mixed-evidence does not allow for a recommendation for chronic use. Cyclobenzaprine is a skeletal muscle relaxant and a central nervous system depressant with similar effects to tricyclic antidepressants (e.g. amitriptyline). Cyclobenzaprine is more effective than placebo in the management of back pain, although the effect is modest and comes at the price of adverse effects." Per p41 of the MTUS guidelines, the effect is greatest in the first 4 days of treatment,

suggesting that shorter courses may be better. Treatment is recommended for the treatment of acute spasm limited to a maximum of 2-3 weeks. UDS that evaluate for cyclobenzaprine can provide additional data on whether the injured worker is compliant, however in this case there is no UDS testing for cyclobenzaprine. The documentation submitted for review indicates that the injured worker has been using this medication since at least 3/2015. There is no documentation of the patients' specific functional level or percent improvement with treatment with cyclobenzaprine. As it is recommended only for short-term use, medical necessity cannot be affirmed.

Ambien 10mg: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter, Insomnia treatment.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Zolpidem (ambien).

Decision rationale: The MTUS is silent on the treatment of insomnia. With regard to Ambien, the ODG guidelines state "Zolpidem is a prescription short-acting non-benzodiazepine hypnotic, which is approved for the short-term (usually two to six weeks) treatment of insomnia. Proper sleep hygiene is critical to the individual with chronic pain and often is hard to obtain. Various medications may provide short-term benefit. While sleeping pills, so-called minor tranquilizers, and anti-anxiety agents are commonly prescribed in chronic pain, pain specialists rarely, if ever, recommend them for long-term use. They can be habit-forming, and they may impair function and memory more than opioid pain relievers. There is also concern that they may increase pain and depression over the long-term." The documentation submitted for review does not contain information regarding sleep onset, sleep maintenance, and sleep quality and next-day functioning. It was not noted whether simple sleep hygiene methods were tried and failed. The request is not medically necessary.