

Case Number:	CM15-0115667		
Date Assigned:	06/23/2015	Date of Injury:	07/07/2009
Decision Date:	07/22/2015	UR Denial Date:	06/01/2015
Priority:	Standard	Application Received:	06/15/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 38-year-old male, who sustained an industrial injury on 7/7/2009. He reported back pain. Diagnoses have included L5-S1 large disc herniation with bilateral radiculopathy. Treatment to date has included L5-S1 discectomy and fusion (5/13/2015), aquatic therapy, epidural steroid injections and medication. According to the progress report dated 5/27/2015, the injured worker reported that his recent back surgery relieved 50% of his sciatica pain in the legs. He was able to discontinue his Dilaudid. He was using Norco, Tramadol, Gabapentin and Flexeril for pain and spasms. Lumbar spine exam revealed a stapled, healing, midline scar. There was palpable spasm and positive tenderness with multiple trigger points. Gait was mildly antalgic on the left. Authorization was requested for Norco, Gabapentin and Flexeril.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

150 Norco 10/325mg: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for low back pain "except for short use for severe cases, not to exceed 2 weeks". The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life". The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. Weaning has been recommended previously. As such, the request for 150 Norco 10/325mg is not medically necessary.

90 Gabapentin 600mg: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-Epilepsy Drugs Page(s): 16-22. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, Anti-epilepsy drugs (AEDs) for pain, Gabapentin (Neurontin ½).

Decision rationale: The MTUS considers Gabapentin as a first-line treatment for neuropathic pain and effective for the treatment of spinal cord injury, lumbar spinal stenosis, and post op pain. MTUS also recommends a trial of Gabapentin for complex regional pain syndrome. ODG states "Recommended Trial Period: One recommendation for an adequate trial with Gabapentin is three to eight weeks for titration, then one to two weeks at maximum tolerated dosage. (Dworkin, 2003) The patient should be asked at each visit as to whether there has been a change in pain or function. Current consensus based treatment algorithms for diabetic neuropathy suggests that if inadequate control of pain is found, a switch to another first-line drug is recommended". Additionally, ODG states that Gabapentin "has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain". The treating physician has not provided documentation of objective functional improvement with the use of this medication. As such, the request for 90 Gabapentin 600mg is not medically necessary.

30 Flexeril 7.5mg: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants (for pain).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine, Medications for chronic pain, Antispasmodics Page(s): 41-42, 60-61, 64-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Cyclobenzaprine (Flexeril ½) and Other Medical Treatment Guidelines UpToDate, Flexeril.

Decision rationale: MTUS Chronic Pain Medical Treatment states for Cyclobenzaprine, "Recommended as an option, using a short course of therapy. The effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. (Browning, 2001) Treatment should be brief. "Additionally, MTUS outlines that "Relief of pain with the use of medications is generally temporary, and measures of the lasting benefit from this modality should include evaluating the effect of pain relief in relationship to improvements in function and increased activity. Before prescribing any medication for pain the following should occur: (1) determine the aim of use of the medication; (2) determine the potential benefits and adverse effects; (3) determine the patient's preference. Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005)" Uptodate "flexeril" also recommends "Do not use longer than 2-3 weeks". Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine. ODG states regarding cyclobenzaprine, "Recommended as an option, using a short course of therapy. The addition of cyclobenzaprine to other agents is not recommended". Other pain medications are being requested, along with cyclobenzaprine, which ODG recommends against. Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine. The medical documentation provided indicates this patient has previously utilized Flexeril without documentation of objective functional improvement. As such, the request for 30 Flexeril 7.5mg is not medically necessary.