

Case Number:	CM15-0116807		
Date Assigned:	06/25/2015	Date of Injury:	02/15/2012
Decision Date:	07/23/2015	UR Denial Date:	06/05/2015
Priority:	Standard	Application Received:	06/17/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 59-year-old male who sustained an industrial injury on 02/15/12. He reports low back pain and constant neck pain. Diagnoses include cervicgia and lumbago. Treatments to date include medial branch block, physical therapy, chiropractic treatment, epidural steroid injections, and pain medication. In a progress note dated 05/12/15, the treating provider reports straight leg raise, Patrick's, and Spurling's tests were negative. Facet loading was positive; there was tenderness to palpation over the cervical paraspinal musculature, and lumbar paraspinal musculature. The injured worker had undergone medial branch block on April 30, 2015 with positive results and significant improvement. Treatment recommendation is for radiofrequency ablation of bilateral L4-5, and L5-S1, and continuation of Tramadol 50 mg for breakthrough pain. Date of Utilization Review: 06/05/15.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Bilateral lumbar radiofrequency rhizotomy at L4-L5 and L5-S1 with fluoroscopy: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 300-301.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 287-315. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Low Back - Lumbar & Thoracic (Acute & Chronic), Facet joint radiofrequency neurotomy.

Decision rationale: ODG states, "Criteria for use of facet joint radiofrequency neurotomy: (1) Treatment requires a diagnosis of facet joint pain using a medial branch block as described above. See Facet joint diagnostic blocks (injections). (2) While repeat neurotomies may be required, they should not occur at an interval of less than 6 months from the first procedure. A neurotomy should not be repeated unless duration of relief from the first procedure is documented for at least 12 weeks at 50% relief. The current literature does not support that the procedure is successful without sustained pain relief (generally of at least 6 months duration). No more than 3 procedures should be performed in a year's period. (3) Approval of repeat neurotomies depends on variables such as evidence of adequate diagnostic blocks, documented improvement in VAS score, decreased medications and documented improvement in function. (4) No more than two joint levels are to be performed at one time. (5) If different regions require neural blockade, these should be performed at intervals of no sooner than one week, and preferably 2 weeks for most blocks. (6) There should be evidence of a formal plan of additional evidence-based conservative care in addition to facet joint therapy." ODG and MD Guidelines agree that: "One diagnostic facet joint injection may be recommended for patients with chronic low back pain that is significantly exacerbated by extension and rotation or associated with lumbar rigidity and not alleviated with other conservative treatments (e.g., NSAIDs, aerobic exercise, other exercise, manipulation) in order to determine whether specific interventions targeting the facet joint are recommended. If after the initial block/blocks are given (see "Diagnostic Phase" above) and found to produce pain relief of at least 50-70% pain relief for at least 6-8 weeks, additional blocks may be supported." The medical documentation provided indicate this patient had a previous medial branch block. Although the physician documents that the patient had relief from the injection, there is no documentation that meets the requirements as described above. As such, the request for bilateral lumbar radiofrequency rhizotomy at L4-L5 and L5-S1 with fluoroscopy is not medically necessary.

Tramadol 50mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Tramadol, Ultram Page(s): 74-96, 113, 123. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic) - Medications for acute pain (analgesics), Tramadol (Ultram).

Decision rationale: Tramadol is classified as a central acting synthetic opioids. MTUS states regarding tramadol that "A therapeutic trial of opioids should not be employed until the patient has failed a trial of non-opioid analgesics. Before initiating therapy, the patient should set goals, and the continued use of opioids should be contingent on meeting these goals." ODG further states, "Tramadol is not recommended as a first-line oral analgesic because of its inferior

efficacy to a combination of Hydrocodone/acetaminophen." The treating physician did not provide sufficient documentation that the patient has failed a trial of non-opioid analgesics at the time of prescription or in subsequent medical notes. Additionally, no documentation was provided which discussed the setting of goals for the use of tramadol prior to the initiation of this medication. MTUS states that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for Tramadol 50mg #60 is not medically necessary.