

Case Number:	CM15-0117536		
Date Assigned:	06/25/2015	Date of Injury:	03/21/2012
Decision Date:	07/27/2015	UR Denial Date:	06/04/2015
Priority:	Standard	Application Received:	06/18/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 39-year-old male, who sustained an industrial injury on 3/21/12. The injured worker has complaints of leg burning and pain. The documentation noted that the injured worker was paxil for depression and anxiety. The diagnoses have included T11 compression fraction; back pain and depression. Treatment to date has included oxycontin; norco and neurontin. The request was for gralise 600mg #30; norco 10/325mg #180 and oxycontin 10mg #60. Several documents within the submitted medical records are difficult to decipher.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Gralise 600mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Gabapentin Page(s): 49.

Decision rationale: According to MTUS guidelines, "Gabapentin is an anti-epilepsy drug (AEDs - also referred to as anti-convulsants), which has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain. " There was no documentation that the patient is suffering from neuropathic pain including diabetic neuropathic pain or post-herpetic neuralgia condition.

Therefore, the prescription of Gralise 600mg #30 is not medically necessary.

Norco 10/325mg #180: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, Norco (Hydrocodone/Acetaminophen) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: "(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug- related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. "According to the patient's file, there is no objective documentation of pain and functional improvement to justify continuous use of Norco. Norco was used for longtime without documentation of functional improvement or evidence of return to work or improvement of activity of daily living. Therefore, the prescription of Norco 10/325mg #180 is not medically necessary.

Oxycontin 10mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, Oxycodone as well as other short acting opioids are indicated for intermittent or breakthrough pain (page 75). It can be used in acute post operative pain. It is not recommended for chronic pain of longterm use as prescribe in this case. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: "(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain;

intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. "There is no clear documentation for the need for continuous use of Oxycontin. There is no documentation for pain and functional improvement with previous use of Oxycontin. There is no documentation of compliance of the patient with his medications. Based on the above, the prescription of OxyContin 10mg #60 is not medically necessary.