

Case Number:	CM15-0117614		
Date Assigned:	06/25/2015	Date of Injury:	05/30/1997
Decision Date:	07/24/2015	UR Denial Date:	06/09/2015
Priority:	Standard	Application Received:	06/18/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California
 Certification(s)/Specialty: Physical Medicine & Rehabilitation

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 47 year old female, who sustained an industrial injury on 5/30/1997. Diagnoses have included cervical spondylosis with acute exacerbation, lumbar myofascial pain and status post bilateral carpal tunnel releases. Treatment to date has included trigger point injections and medication. According to the progress report dated 5/14/2015, the injured worker complained of neck pain and stiffness. She also complained of intermittent back pain. Objective findings revealed tenderness in the posterior cervical and bilateral trapezial musculature. There was tenderness in the lower lumbar paravertebral musculature. Authorization was requested for Ultram, Prilosec and a urine toxicology screen.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Ultram 50mg #60 with 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Criteria for use.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

Decision rationale: Per the MTUS Guidelines, cite opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury of 1997 without acute flare, new injury, or progressive deterioration. The Ultram 50mg #60 with 2 refills is not medically necessary and appropriate.

Prilosec 20mg #30 with 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI Symptoms and Cardiovascular risk, Pages 68-69.

Decision rationale: Proton pump inhibitor (PPI) medication is for treatment of the problems associated with active gastric ulcers, erosive esophagitis, Barrett's esophagitis, or in patients with pathologic hypersecretion diseases. Although preventive treatment is effective for the mentioned diagnosis, studies suggest; however, nearly half of PPI prescriptions are used for unapproved or no indications. Per MTUS Chronic Pain Treatment Guidelines, the patient does not meet criteria for Omeprazole (Prilosec) namely reserved for patients with history of prior GI bleeding, the elderly (over 65 years), diabetics, and chronic cigarette smokers. Long term use of PPIs have potential increased risks of B12 deficiency; iron deficiency; hypomagnesemia; susceptibility to pneumonia, enteric infections, fractures, hypergastrinemia and cancer, and cardiovascular effects of myocardial infarction (MI). In the elderly, studies have demonstrated increased risk for Clostridium difficile infection, bone loss, and fractures from long-term use of PPIs. Given treatment criteria outweighing risk factors, if a PPI is to be used, omeprazole (Prilosec), lansoprazole (Prevacid), and esomeprazole (Nexium) are to be considered over second-line therapy of other PPIs such as pantoprazole (Protonix), dexlansoprazole (Dexilant), and rabeprazole (Aciphex). Submitted reports have not described or provided any GI diagnosis that meets the criteria to indicate medical treatment. Review of the records show no documentation of any history, symptoms, or GI diagnosis to warrant this medication. The Prilosec 20mg #30 with 2 refills is not medically necessary and appropriate.

One urine toxicology screen: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, steps to avoid misuse/addiction.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Drug Testing, page 43.

Decision rationale: Per MTUS Guidelines, urine drug screening is recommended as an option before a therapeutic trial of opioids and for on-going management to differentiate issues of abuse, addiction, misuse, or poor pain control; none of which apply to this patient who has been prescribed long-term opioid this chronic injury of 1997. Presented medical reports from the provider have unchanged chronic severe pain symptoms with unchanged clinical findings of restricted range and tenderness without acute new deficits or red-flag condition changes. Treatment plan remains unchanged with continued medication refills without change in dosing or prescription for chronic pain. There is no report of aberrant behaviors, illicit drug use, and report of acute injury or change in clinical findings or risk factors to support frequent UDS. Documented abuse, misuse, poor pain control, history of unexpected positive results for a non-prescribed scheduled drug or illicit drug or history of negative results for prescribed medications may warrant UDS and place the patient in a higher risk level; however, none are provided. The One urine toxicology screen is not medically necessary and appropriate.