

Case Number:	CM15-0119404		
Date Assigned:	06/29/2015	Date of Injury:	05/26/2003
Decision Date:	07/29/2015	UR Denial Date:	05/26/2015
Priority:	Standard	Application Received:	06/20/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 56 year old female, who sustained an industrial injury on 5/26/2003. Diagnoses include lumbago, right tibial plateau fracture status post open reduction internal fixation (ORIF) with right knee plating (2003), left knee meniscus injury status-post surgical intervention (2003), and chronic low back pain with radiculopathy verified by EMG (electromyography)/NCV (nerve conduction studies), right worse than left. Treatment to date has included diagnostics, surgical intervention, and conservative care including therapy, medications and injections. Per the Primary Treating Physician's Progress Report dated 5/15/2015 the injured worker reported lower back pain and left knee pain. Physical examination revealed positive straight leg raise bilaterally at about 45 degrees of extension, right worse than left. She does have some pain in the lower extremities. There is radiculopathy from the L5 dermatomes bilaterally but more of the pain is in her big toe, right side greater than left. Physical examination of the knees revealed crepitus with some significant pain. She has been evaluated by the surgeon who feels that there is no surgery needed maybe just possible therapy. The plan of care included continuation of prescribed medications and physical therapy. Authorization was requested for 12 sessions of physical therapy for the knee.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

12 Physical Therapy sessions for the knee: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks-Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2): 8-10 visits over 4 weeks-Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The requested amount of physical therapy is in excess of California chronic pain medical treatment guidelines. There is no objective explanation why the patient would need excess physical therapy and not be transitioned to active self-directed physical medicine. The request is not medically necessary.