

Case Number:	CM15-0120543		
Date Assigned:	07/01/2015	Date of Injury:	12/20/2011
Decision Date:	09/21/2015	UR Denial Date:	05/21/2015
Priority:	Standard	Application Received:	06/22/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Indiana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 44-year-old male, who sustained an industrial injury on 12/20/2011. He reported low back pain while offloading rail cars. The injured worker was diagnosed as having cervical pain, cervical radiculopathy, cervical sprain/strain, thoracic muscle spasm, thoracic pain, thoracic sprain/strain, lumbar sprain/strain, right shoulder muscle spasm, right shoulder pain, and right shoulder sprain/strain. Treatment to date has included diagnostics, physical therapy, chiropractic, epidural steroid injections, lumbar spinal surgery in 2012, and medications. Per the most recent PR2 report (10/13/2014), the injured worker complains of pain to his cervical spine, thoracic spine, lumbar spine, and right shoulder. It was also noted that he was having right knee pain lately. His medications included Naproxen, Tizanidine, Hydrocodone/APAP, and compounded medication creams. He underwent urine screen to rule out toxicity. The results were not noted. His work status was not documented. An updated progress report regarding the current recommendations was not noted.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Retrospective request for Pantoprazole 20mg #60 (DOS: 05/12/15): Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids; GI risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: MTUS states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e. g. , NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1. 44)." ODG states, "If a PPI is used, omeprazole OTC tablets or lansoprazole 24HR OTC are recommended for an equivalent clinical efficacy and significant cost savings. Products in this drug class have demonstrated equivalent clinical efficacy and safety at comparable doses, including esomeprazole (Nexium), lansoprazole (Prevacid), omeprazole (Prilosec), pantoprazole (Protonix), dexlansoprazole (Dexilant), and rabeprazole (Aciphex). (Shi, 2008) A trial of omeprazole or lansoprazole is recommended before Nexium therapy. The other PPIs, Protonix, Dexilant, and Aciphex, should also be second-line. According to the latest AHRQ Comparative Effectiveness Research, all of the commercially available PPIs appeared to be similarly effective. (AHRQ, 2011)" The patient does not meet the age recommendations for increased GI risk. The medical documents provided establish the patient has experienced GI discomfort, but is nonspecific and does not indicate history of peptic ulcer, GI bleeding or perforation. Medical records do not indicate that the patient is on ASA, corticosteroids, and/or an anticoagulant; or high dose/multiple NSAID. Additionally per guidelines, Pantoprazole is considered second line therapy and the treating physician has not provided detailed documentation of a failed trial of omeprazole and/or lansoprazole. As such, the request for Pantoprazole is not medically necessary.

Retrospective request for Diclofenac 100mg #60 (DOS: 05/12/15): Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Diclofenac.

Decision rationale: MTUS specifies four recommendations regarding NSAID use: 1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. 2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP. 3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A

Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. 4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. The treating physician does not document failure of primary (Tylenol) treatment. Importantly, ODG also states that diclofenac is "Not recommended as first line due to increased risk profile . . . If using diclofenac then consider discontinuing as it should only be used for the shortest duration possible in the lowest effective dose due to reported serious adverse events". Medical documents indicate that the patient has been on diclofenac for at least several months, which given the treatment history does not appear to be the shortest duration possible. As such, the request is not medically necessary.

Retrospective request for Cyclobenzaprine 7.5mg #90 (DOS: 05/12/15): Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-spasmodics, cyclobenzaprine Page(s): 41-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Cyclobenzaprine (Flexeril®).

Decision rationale: MTUS Chronic Pain Medical Treatment states for Cyclobenzaprine, "Recommended as an option, using a short course of therapy. . . The effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. (Browning, 2001) Treatment should be brief." The medical documents indicate that patient is far in excess of the initial treatment window and period. Additionally, MTUS outlines that "Relief of pain with the use of medications is generally temporary, and measures of the lasting benefit from this modality should include evaluating the effect of pain relief in relationship to improvements in function and increased activity. Before prescribing any medication for pain the following should occur: (1) determine the aim of use of the medication; (2) determine the potential benefits and adverse effects; (3) determine the patient's preference. Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005)" Uptodate "flexeril" also recommends "Do not use longer than 2-3 weeks". Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine. ODG states regarding cyclobenzaprine, "Recommended as an option, using a short course of therapy . . . The addition of cyclobenzaprine to other agents is not recommended". Several other pain medications are being requested, along with cyclobenzaprine, which ODG recommends against. As such, the request is not medically necessary.

Retrospective request for Norco 10/325mg #120 (DOS: 05/12/15): Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for neck and low back pain "except for short use for severe cases, not to exceed 2 weeks". The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life". The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for Norco 325/10mg # 120 is not medically necessary.

Retrospective request for urine toxicology screen (DOS: 05/12/15): Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids; abuse Page(s): 74-109. Decision based on Non-MTUS Citation University of Michigan Health System Guidelines for Clinical Care: Managing Chronic Non-terminal Pain, Including Prescribing Controlled Substances (May 2009), pg 32 Established Patients Using a Controlled Substance.

Decision rationale: MTUS states that use of urine drug screening for illegal drugs should be considered before therapeutic trial of opioids are initiated. Additionally, "Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion) would indicate need for urine drug screening. There is insufficient documentation provided to suggest issues of abuse, addiction, or poor pain control by the treating physician. University of Michigan Health System Guidelines for Clinical Care: Managing Chronic Non-terminal Pain, Including Prescribing Controlled Substances (May 2009) recommends for stable patients without red flags "twice yearly urine drug screening for all chronic non-malignant pain patients receiving opioids, once during January-June and another July-December". The patient has been on chronic opioid therapy. The treating physician has not indicated why a urine drug screen is necessary at this time and has provided no evidence of red flags. As such, the request is not medically necessary.

Retrospective request for confirmations for medication and monitoring and specimen collections and handling (DOS: 5/12/15): Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids; abuse Page(s): 74-109. Decision based on Non-MTUS Citation University of Michigan Health System Guidelines for Clinical Care: Managing Chronic Non-terminal Pain, Including Prescribing Controlled Substances (May 2009), pg 32 Established Patients Using a Controlled Substance.

Decision rationale: MTUS states that use of urine drug screening for illegal drugs should be considered before therapeutic trial of opioids are initiated. Additionally, "Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion) would indicate need for urine drug screening. There is insufficient documentation provided to suggest issues of abuse, addiction, or poor pain control by the treating physician. University of Michigan Health System Guidelines for Clinical Care: Managing Chronic Non-terminal Pain, Including Prescribing Controlled Substances (May 2009) recommends for stable patients without red flags "twice yearly urine drug screening for all chronic non-malignant pain patients receiving opioids, once during January-June and another July-December". The patient has been on chronic opioid therapy. The treating physician has not indicated why a urine drug screen is necessary at this time and has provided no evidence of red flags. Thus, the urinalysis is not medically necessary, and by extension, this request is not medically necessary.