

<b>Case Number:</b>	CM15-0122074		
<b>Date Assigned:</b>	07/06/2015	<b>Date of Injury:</b>	09/02/2011
<b>Decision Date:</b>	07/31/2015	<b>UR Denial Date:</b>	06/10/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	06/24/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: North Carolina  
 Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 61-year-old male, who sustained an industrial injury on 09/02/2011. Medical records provided by the treating physician did not indicate the injured worker's mechanism of injury. The injured worker was diagnosed as having degeneration of the lumbar or lumbosacral intervertebral disc, displacement of the lumbar intervertebral disc without myelopathy, other symptoms referable to the back, lumbago, lumbar radiculopathy, low back strain, and hip pain. Treatment and diagnostic studies to date has included use of ice, use of heat, home exercise program, medication regimen, and magnetic resonance imaging of the lumbar spine. In a progress note dated 05/27/2015, the treating physician reports complaints of chronic low back pain. Examination reveals mild tenderness to the lumbosacral spine at lumbar four through sacral one, positive straight leg raise, restricted lumbar range of motion, and hypoesthesia. The injured worker's current medication regimen included Norco, Prilosec, Fish Oil, Lisinopril, Pravastatin, and Paroxetine. The injured worker's pain level is rated a 5 out of 10 with and without use of his medication regimen. The treating physician noted that use of the injured worker's medication regimen along with activity restriction and rest has made the pain manageable to allow the injured worker to complete activities of daily living. The treating physician requested Norco 10/325mg with a quantity of 60 with the treating physician noting that the injured worker's medication regimen keeps the injured worker's pain level manageable allowing the injured worker to complete activities of daily living.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Norco 10/325mg #60:** Overturned

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

**Decision rationale:** The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions are from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work; (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. These criteria are met in the provided clinical documentation and the request is medically necessary.