

Case Number:	CM15-0123152		
Date Assigned:	07/07/2015	Date of Injury:	07/09/2013
Decision Date:	08/13/2015	UR Denial Date:	05/26/2015
Priority:	Standard	Application Received:	06/25/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 60-year-old male who sustained an industrial injury on 07/09/2013. There was no mechanism of injury documented. The injured worker was diagnosed with lumbar intervertebral disc with myelopathy, lumbar sprain/strain and insomnia. No surgical interventions were documented. Treatment to date has included diagnostic testing, acupuncture therapy, chiropractic therapy and medications. According to the primary treating physician's progress report on May 5, 2015, the injured worker continues to experience low back pain rated at 8/10 on the pain scale with sleep disturbance. Examination of the lumbar spine demonstrated decreased range of motion with flexion at 50 degrees, extension at 15 degrees, and bilateral lateral bending at 15 degrees each. Bilateral straight leg raise caused pain. Current medications are listed as Tramadol 15mg and Naproxen. Treatment plan consists of continuing with medication regimen and the current request for continued chiropractic therapy twice a week for 3 weeks for the lumbar spine, continued acupuncture therapy twice a week for 3 weeks for the lumbar spine and a urinalysis for drug screening.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Chiropractic treatment for lumbar spine: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Manual therapy & manipulation.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines manual manipulation Page(s): 58-59.

Decision rationale: The California chronic pain medical guidelines section on manual manipulation states: Recommended for chronic pain if caused by musculoskeletal conditions. Manual Therapy is widely used in the treatment of musculoskeletal pain. The intended goal or effect of Manual Medicine is the achievement of positive symptomatic or objective measurable gains in functional improvement that facilitate progression in the patient's therapeutic exercise program and return to productive activities. Manipulation is manual therapy that moves a joint beyond the physiologic range-of-motion but not beyond the anatomic range-of-motion. Low back: Recommended as an option. Therapeutic care & Trial of 6 visits over 2 weeks, with evidence of objective functional improvement, total of up to 18 visits over 6-8 weeks. Elective/maintenance care & Not medically necessary. Recurrences/flare-ups, Need to reevaluate treatment success, if RTW achieved then 1-2 visits every 4-6 months. Ankle & Foot: Not recommended. Carpal tunnel syndrome: Not recommended. Forearm, Wrist, & Hand: Not recommended. Knee: Not recommended. Treatment Parameters from state guidelines: a. Time to produce effect: 4 to 6 treatments Manual manipulation is recommended form of treatment for chronic pain. However, the requested amount of therapy sessions is in excess of the recommendations per the California MTUS. The California MTUS states there should be not more than 6 visits over 2 weeks and documented evidence of functional improvement before continuation of therapy. The patient has had chiropractic care without documented objective measurements of significant improvement in pain and function. Therefore, the request is denied for additional therapy. This request is not medically necessary.

Urinalysis: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, ongoing use.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a

framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. The California MTUS does recommend urine drug screens as part of the criteria for ongoing use of opioids. The patient was on opioids at the time of request and therefore the request is medically necessary.

Acupuncture for lumbar spine, 2x3: Upheld

Claims Administrator guideline: Decision based on MTUS Acupuncture Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Acupuncture Treatment Guidelines.

Decision rationale: The California chronic pain medical treatment guidelines section on acupuncture states: Acupuncture is used as an option when pain medication is reduced or not tolerated, it may be used as an adjunct to physical rehabilitation and/or surgical intervention to hasten functional recovery. It is the insertion and removal of filiform needles to stimulate acupoints (acupuncture points). Needles may be inserted, manipulated, and retained for a period of time. Acupuncture can be used to reduce pain, reduce inflammation, increase blood flow, increase range of motion, decrease the side effect of medication-induced nausea, promote relaxation in an anxious patient, and reduce muscle spasm. Time to produce functional improvement is 3-6 treatments and frequency is 1-3 times per week. The patient has had acupuncture without documented objective measurements of significant improvement in pain and function. Therefore, additional sessions are not medically necessary.