

Case Number:	CM15-0125670		
Date Assigned:	07/10/2015	Date of Injury:	04/03/2000
Decision Date:	08/06/2015	UR Denial Date:	06/03/2015
Priority:	Standard	Application Received:	06/29/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is 53 year old male who sustained an industrial injury 4/03/2000. Diagnoses are spinal enthesopathy, cervicalgia, spasm of muscle, and lumbago. In a progress report date 4/13/15, a treating physician notes the injured worker has ongoing pain in the lumbar area with pain levels at 5-6/10 on average. When it travels to the leg, it reaches 10/10. Neck pain is constant at 4-5/10, left arm pain is at 2-3/10, and he has pain around the shoulder blades as well with muscle cramping and spasms. Pain medication lessens pain and allows for increased function in activities of daily living. Medications are Flexeril, Celebrex, Norco, and AmbienCR. He has evidence on an MRI of degenerative changes and prior surgery at the L4,5 area. The spine has mild tenderness in the lumbar area with good range of motion. There is tenderness of the thoracic paraspinal muscles and trap area. Deep tendon reflexes are decreased slightly in the right patellar and bilateral achilles. The lumbar spine xray done 4/13/15 demonstrates further degenerative changes at L4, L5 and L5-S1 with narrowing of the disc space. This is a pronounced loss of disc height at L5, S1. Compared to prior studies, further degenerative changes and lipping is noted. The requested treatment is an MRI of the lumbar spine with contrast.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MRI with contrast: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG): Low Back - Lumbar & Thoracic (Acute & Chronic) Chapter (Online Version).

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304.

Decision rationale: The ACOEM chapter on low back complaints and special diagnostic studies states: Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computed tomography [CT] for bony structures). Relying solely on imaging studies to evaluate the source of low back and related symptoms carries a significant risk of diagnostic confusion (false positive test results) because of the possibility of identifying a finding that was present before symptoms began and therefore has no temporal association with the symptoms. Techniques vary in their abilities to define abnormalities (Table 12-7). Imaging studies should be reserved for cases in which surgery is considered or red-flag diagnoses are being evaluated. Because the overall false-positive rate is 30% for imaging studies in patients over age 30 who do not have symptoms, the risk of diagnostic confusion is great. There is no recorded presence of emerging red flags on the physical exam. There is evidence of nerve compromise on physical exam but there is not mention of consideration for surgery or complete failure of conservative therapy. There are no significant changes in exam since previous MRI. For these reasons, criteria for imaging as defined above per the ACOEM have not been met. Therefore the request is not medically necessary.