

Case Number:	CM15-0126124		
Date Assigned:	07/10/2015	Date of Injury:	11/26/2003
Decision Date:	08/06/2015	UR Denial Date:	06/19/2015
Priority:	Standard	Application Received:	06/30/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51-year-old male who sustained an industrial injury on 11/26/2003. Mechanism of injury occurred when he tried to lift construction material and felt a pop in his lower back and shooting pain in his buttocks and legs, and fell to the ground. Diagnoses include chronic pain syndrome, failed lumbar back surgery, post laminectomy pain, lumbar back pain with bilateral radiculopathy, degenerative lumbar spine disc disease, history of arthrodesis L4-L5, L5-S1, anterior and posterior fusion, depression, insomnia, anxiety and multiple back operations. Treatment to date has included diagnostic studies, medications, intrathecal pain pump; he uses a cane or wheel chair, 5 lumbar surgeries, 3 thoracic surgeries, and one cervical surgery, Cognitive Behavioral Therapy, spinal cord stimulator, and physical therapy. Current medications include Norco, Constulose, Lexapro, Lunesta, Imipramine, Provigil, and intrathecal Dilaudid. A physician progress note dated 06/11/2015 documents the injured worker has pain in the head, arms, legs, neck, left shoulder, buttocks, elbows, hips, hands, knees, abdomen, low back, ankles/feet and groin. He rests or reclines most of the day and is not up and out of bed daily. His intrathecal pump was refilled today. His pain level is high. Following one of his surgeries in 2012 he developed urinary incontinence. He has chronic mild lower abdominal pain and onyx has a bowel movement about every 4-5 days. His average pain is 6 out of 10 and his worst pain is 8 out of 10 with one being the least pain and 10 being the worst pain. Pain and spasticity is constant. Treatment requested is for Movantik 12. 5mg #60 with 2 refills and Norco 10/325mg #180.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #180: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): s 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs (Passik, 2000). (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997), (Wisconsin, 2004) (Warfield, 2004). The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

Movantik 12. 5mg #60 with 2 refills: Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 77.

Decision rationale: The California chronic pain medical treatment guidelines section on opioid therapy states: (a) Intermittent pain: Start with a short-acting opioid trying one medication at a time. (b) Continuous pain: extended-release opioids are recommended. Patients on this modality may require a dose of "rescue" opioids. The need for extra opioid can be a guide to determine the sustained release dose required. (c) Only change 1 drug at a time. (d) Prophylactic treatment of constipation should be initiated. The patient is currently on opioid therapy. The use of constipation measures is advised per the California MTUS. The requested medication is used in the treatment of constipation. Therefore, the request is medically necessary.