

<b>Case Number:</b>	CM15-0126211		
<b>Date Assigned:</b>	07/10/2015	<b>Date of Injury:</b>	01/30/2003
<b>Decision Date:</b>	08/06/2015	<b>UR Denial Date:</b>	06/17/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	06/30/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51-year-old male, who sustained an industrial injury on 1/30/2003. The mechanism of injury is unknown. The injured worker was diagnosed as having chronic pain syndrome and lumbar post laminectomy syndrome. There is no record of a recent diagnostic study. Treatment to date has included therapy and medication management. In a progress note dated 6/10/2015, the injured worker complains of muscle aches, joint pains and low back pain, rated 10/10 without medications and 4/10 with medications. Physical examination showed lumbar paraspinal tenderness and decreased sensation in the left lower extremity. The treating physician is requesting Clonazepam 2 mg, #30, Dilaudid 8 mg, #150 and Methadone 10 mg, #180.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Clonazepam 2mg tablet #30:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

**Decision rationale:** The MTUS/Chronic Pain Medical Treatment Guidelines comment on the use of benzodiazepines, including Clonazepam, as a treatment modality. Benzodiazepines are not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks. Their range of action includes sedative/hypnotic, anxiolytic, anticonvulsant, and muscle relaxant. Chronic benzodiazepines are the treatment of choice in very few conditions. Tolerance to hypnotic effects develops rapidly. Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. A more appropriate treatment for anxiety disorder is an antidepressant. Tolerance to anticonvulsant and muscle relaxant effects occurs within weeks. In this case, the records indicate that the use of Clonazepam has extended well-beyond the 4-week limit as described in the above cited MTUS guidelines. There is insufficient justification provided in support of long-term use. For this reason, Clonazepam is not considered as medically necessary. In the Utilization Review process, a limited supply of Clonazepam was provided to allow for weaning. This action is consistent with the above cited guidelines. The request is not medically necessary.

**Dilaudid 8mg tablet #150:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 76-78, 80.

**Decision rationale:** The MTUS/Chronic Pain Medical Treatment Guidelines comment on the long-term use of opioids, including Dilaudid. These guidelines have established criteria of the use of opioids for the ongoing management of pain. Actions should include: prescriptions from a single practitioner and from a single pharmacy. The lowest possible dose should be prescribed to improve pain and function. There should be an ongoing review and documentation of pain relief, functional status, appropriate medication use and side effects. Pain assessment should include: current pain, the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. There should be evidence of documentation of the 4 A's for Ongoing Monitoring. These four domains include: pain relief, side effects, physical and psychological functioning, and the occurrence of any potentially aberrant drug-related behaviors. Further, there should be consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain that does not improve on opioids in 3 months. There should be consideration of an addiction medicine consult if there is evidence of substance misuse (Pages 76-78). Finally, the guidelines indicate that for chronic pain, the long-term efficacy of opioids is unclear. Failure to respond to a time-limited course of opioids has led to the suggestion of reassessment and consideration of alternative therapy (Page 80). Based on the review of the medical records, there is insufficient documentation in support of these stated MTUS/Chronic Pain Medical Treatment Guidelines for the ongoing use of opioids. There is insufficient documentation of the 4 A's for Ongoing Monitoring. The treatment course of opioids in this patient has extended well beyond the timeframe required for a reassessment of therapy. In summary, there is insufficient documentation to support the chronic use of an opioid in this patient. Treatment with Dilaudid is not medically necessary. In the Utilization Review process,

a limited supply of Dilaudid was certified to allow for weaning. This action is consistent with the above cited guidelines.

**Methadone 10mg tablet #180:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 76-78, 80.

**Decision rationale:** The MTUS/Chronic Pain Medical Treatment Guidelines comment on the long-term use of opioids, including Methadone. These guidelines have established criteria of the use of opioids for the ongoing management of pain. Actions should include: prescriptions from a single practitioner and from a single pharmacy. The lowest possible dose should be prescribed to improve pain and function. There should be an ongoing review and documentation of pain relief, functional status, appropriate medication use and side effects. Pain assessment should include: current pain, the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. There should be evidence of documentation of the 4 A's for Ongoing Monitoring. These four domains include: pain relief, side effects, physical and psychological functioning, and the occurrence of any potentially aberrant drug-related behaviors. Further, there should be consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain that does not improve on opioids in 3 months. There should be consideration of an addiction medicine consult if there is evidence of substance misuse (Pages 76-78). Finally, the guidelines indicate that for chronic pain, the long-term efficacy of opioids is unclear. Failure to respond to a time-limited course of opioids has led to the suggestion of reassessment and consideration of alternative therapy (Page 80). Based on the review of the medical records, there is insufficient documentation in support of these stated MTUS/Chronic Pain Medical Treatment Guidelines for the ongoing use of opioids. There is insufficient documentation of the 4 A's for Ongoing Monitoring. The treatment course of opioids in this patient has extended well beyond the timeframe required for a reassessment of therapy. In summary, there is insufficient documentation to support the chronic use of an opioid in this patient. Treatment with Methadone is not medically necessary.