

<b>Case Number:</b>	CM15-0126525		
<b>Date Assigned:</b>	07/13/2015	<b>Date of Injury:</b>	02/20/2013
<b>Decision Date:</b>	08/06/2015	<b>UR Denial Date:</b>	06/03/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	06/30/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 57-year-old male with an industrial injury dated 02/20/2013. The injured worker's diagnoses include left knee degenerative joint disease, status post left total knee arthroplasty and left knee arthrofibrosis. Treatment consisted of diagnostic studies, prescribed medications, and periodic follow up visits. In a progress note dated 04/08/2015, the injured worker reported tightness, pain and swelling in the left knee. Objective findings revealed swelling, large effusion and pain primarily over the anterior aspect of the left knee in the tibial region. The treating physician prescribed Norco 7.5/325mg #120 and water therapy 3x6 now under review.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Norco 7.5/325 MG #120:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines opioids  
Page(s): 76-84.

**Decision rationale:** The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids; (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004). The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

**Water Therapy 3x6:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Aqua Therapy.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines aqua therapy Page(s): 22.

**Decision rationale:** The California chronic pain medical treatment guidelines section on aquatic therapy states: Aquatic therapy; Recommended as an optional form of exercise therapy, where available, as an alternative to land based physical therapy. Aquatic therapy (including swimming) can minimize the effects of gravity, so it is specifically recommended where reduced weight bearing is desirable, for example extreme obesity. For recommendations on the number of supervised visits, see Physical medicine. Water exercise improved some components of health-related quality of life, balance, and stair climbing in females with fibromyalgia, but regular exercise and higher intensities may be required to preserve most of these gains. (Tomas-Carus, 2007) There is no indication in the provided documentation that this patient has a condition such as extreme obesity that would preclude the patient from land-based physical therapy. The request for physical therapy is not within the recommended number of session even though the need for aquatic versus land based physical therapy has not been established. For these reasons criteria have not been met for the requested service and it is not medically necessary.