

Case Number:	CM15-0127354		
Date Assigned:	07/14/2015	Date of Injury:	08/12/1998
Decision Date:	08/10/2015	UR Denial Date:	06/26/2015
Priority:	Standard	Application Received:	07/01/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 57-year-old female who sustained an industrial injury on 8/12/1998 resulting in right wrist pain. She was diagnosed with traumatic tear of triangular fibrocartilage complex of right wrist; probable right carpal tunnel syndrome; and, carpal boss of right wrist. Documented treatment has included traumatic tear of triangular fibrocartilage repair and excision of carpal boss. Other treatments are not discussed in the provided documentation. The injured worker continues to complain of wrist pain. The treating physician's plan of care includes 8 sessions of occupational therapy for the right wrist and electromyography and nerve conduction velocity of upper extremity. Work status is not addressed in provided documentation.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Occupational Therapy for the right wrist quantity 8 sessions: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 11 Forearm, Wrist, and Hand Complaints.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks; Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks; Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks; The requested amount of physical therapy meets standards as cited above. The patient has an injury amenable to physical therapy. Therefore the request is certified.

Electromyography/Nerve Conduction Velocity of upper extremity: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 11 Forearm, Wrist, and Hand Complaints.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 173-174.

Decision rationale: The ACOEM chapter on neck and upper back complaints and special diagnostic studies states: Criteria for ordering imaging studies are: Emergence of a red flag Physiologic evidence of tissue insult or neurologic dysfunction failure to progress in a strengthening program intended to avoid surgery. Clarification of the anatomy prior to an invasive procedure. Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained

before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H- reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The assessment may include sensory- evoked potentials (SEPs) if spinal stenosis or spinal cord myelopathy is suspected. If physiologic evidence indicates tissue insult or nerve impairment, consider a discussion with a consultant regarding next steps, including the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computed tomography [CT] for bony structures). Additional studies may be considered to further define problem areas. The recent evidence indicates cervical disk annular tears may be missed on MRIs. The clinical significance of such a finding is unclear, as it may not correlate temporally or anatomically with symptoms. The provided documentation does not show any signs of emergence of red flags or subtle physiologic evidence of tissue insult or neurologic dysfunction. There is no mention of planned invasive procedures. There are no subtle neurologic findings listed on the physical exam. For these reasons criteria for special diagnostic testing has not been met per the ACOEM. Therefore the request is not medically necessary.