

Case Number:	CM15-0128022		
Date Assigned:	07/17/2015	Date of Injury:	04/05/2007
Decision Date:	08/13/2015	UR Denial Date:	06/26/2015
Priority:	Standard	Application Received:	07/02/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51 year old male who sustained an industrial injury on 04/05/07. Initial complaints and diagnoses are not available. Treatments to date include medications and right shoulder surgery. Diagnostic studies include a MRI of the right shoulder. Current complaints include pain in the right shoulder and right wrist. Current diagnoses include right shoulder bursitis, right carpal tunnel syndrome, and right wrist tenosynovitis. In a progress note dated 06/23/15 the treating provider reports the plan of care as medications including Flurbiprofen/baclofen/camphor/menthol/dexamethasone/capsaicin/hyaluronic acid, Norco, as well as home exercises, a MRI of the right shoulder, and electrodiagnostic studies of the bilateral upper extremities. The requested treatments include Norco, Flurbiprofen/baclofen/camphor /menthol/dexamethasone/capsaicin/hyaluronic acid, and electrodiagnostic studies of the bilateral upper extremities.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Flurbiprofen 20% / Baclofen 5% / Camphor 2% / Menthol 2% / Dexamethasone Micro 0.2% / Capsaicin 0.025% / Hyaluronic Acid 0.2% in cream base 240gm quantity: 1:
 Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics, Compounded. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain (Chronic), Topical Analgesics (2015).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines topical analgesics Page(s): 111-113.

Decision rationale: The California chronic pain medical treatment guidelines section on topical analgesics states: Recommended as an option as indicated below. Largely experimental in use with few randomized controlled trials to determine efficacy or safety primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. (Namaka, 2004) These agents are applied locally to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. (Colombo, 2006) Many agents are compounded as monotherapy or in combination for pain control (including NSAIDs, opioids, capsaicin, local anesthetics, antidepressants, glutamate receptor antagonists, α -adrenergic receptor agonist, adenosine, cannabinoids, cholinergic receptor agonists, β -agonists, prostanoids, bradykinin, adenosine triphosphate, biogenic amines, and nerve growth factor). (Argoff, 2006) There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The requested medication contains ingredients (hyaluronic acid), which are not indicated per the California MTUS for topical analgesic use. Therefore the request is not medically necessary.

Norco 10/325mg # 90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring

the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

EMG (Electromyography)/ NCV (Nerve Conduction Velocity) of the bilateral upper extremities: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 178.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 173-174.

Decision rationale: The ACOEM chapter on neck and upper back complaints and special diagnostic studies states: Criteria for ordering imaging studies are: Emergence of a red flag. Physiologic evidence of tissue insult or neurologic dysfunction. Failure to progress in a strengthening program intended to avoid surgery-Clarification of the anatomy prior to an invasive procedure. Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The assessment may include sensory-evoked potentials (SEPs) if spinal stenosis or spinal cord myelopathy is suspected. If physiologic evidence indicates tissue insult or nerve impairment, consider a discussion with a consultant regarding next steps, including the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computer tomography [CT] for bony structures). Additional studies may be considered to further define problem areas. The recent evidence indicates cervical disk annular tears may be missed on MRIs. The clinical significance of such a finding is unclear, as it may not correlate temporally or anatomically with symptoms. The provided documentation does not show any signs of emergence of red flags or physiologic evidence of tissue insult or neurologic dysfunction. There is no mention of planned invasive procedures. There are no subtle neurologic findings listed on the physical exam. For these reasons criteria for special diagnostic testing has not been met per the ACOEM. Therefore the request is not medically necessary.