

Case Number:	CM15-0128115		
Date Assigned:	07/14/2015	Date of Injury:	01/01/2005
Decision Date:	08/12/2015	UR Denial Date:	06/25/2015
Priority:	Standard	Application Received:	07/02/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 62 year old female, who sustained an industrial injury on January 01, 2005. Medical records provided by the treating physician did not indicate the injured worker's mechanism of injury. The injured worker was diagnosed as having shoulder pain, entrapment neuropathy of the upper limb, extremity pain, elbow pain, carpal tunnel syndrome, ulnar neuropathy, lateral epicondylitis, status post right shoulder revision of a rotator cuff repair, status post right shoulder extensive debridement, status post right shoulder arthroscopic revision acromioplasty, and status post right shoulder revision of lateral scar. Treatment and diagnostic studies to date has included above noted procedures, medication regimen, physical therapy, status post right bicipital groove injection under ultrasound guidance, magnetic resonance imaging of the right shoulder, electromyogram with an nerve conduction study of the bilateral upper extremity, magnetic resonance imaging of the right shoulder, and a home exercise program. In a progress note dated May 13, 2015 the treating physician noted pain to the neck, pain to the bilateral upper extremities, and pain to the right shoulder. In a progress note dated June 01, 2015 the examination reveals crepitus to the subacromial space, decreased range of motion to the right shoulder, and decreased strength to the rotator cuff. Physical therapy progress note from March 12, 2015 was noted to be the injured worker's 26th visit and was remarkable for increased crepitus to the shoulder with the biceps more painful and noted a questionable rupture along with the treating physical therapist noting that the injured worker to be progressing, but the documentation did not indicate if the injured worker experienced any functional improvement

from the prior physical therapy. The treating physician requested six sessions of physical therapy to the right shoulder with the treating physician noting that the injured worker was not progressing with her home exercise program and was starting to have a loss of range of motion due to lack of further physical therapy.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Physical therapy for the right shoulder, QTY: 6: Upheld

Claims Administrator guideline: Decision based on MTUS Postsurgical Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The patient has already completed physical therapy. The request is in excess of these recommendations per the California MTUS. There is no objective reason why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions in the provided clinical documentation. Therefore, the request is not medically necessary.

