

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM15-0128617 | | |
| Date Assigned: | 07/15/2015 | Date of Injury: | 05/28/2001 |
| Decision Date: | 09/21/2015 | UR Denial Date: | 06/17/2015 |
| Priority: | Standard | Application Received: | 07/02/2015 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker (IW) is a 59-year-old male who sustained an industrial injury on 05/28/2001. Diagnoses include cervical disc displacement; lumbar disc displacement; symptoms of anxiety and depression; carpal tunnel syndrome; and cervical disc degeneration. Treatment to date has included medication. According to the progress notes dated 5/22/15, the IW reported pain and discomfort in the cervical spine and upper extremity region. Daily living activities aggravated the pain and the pain made sleep difficult. She also complained of stomach pain due to prolonged medication use. On examination, muscle tightness and spasms were present in the trapezius, sternocleidomastoid and straps, bilaterally. Range of motion of the cervical spine was: forward flexion 50 degrees, extension 50 degrees, rotation right and left 65 degrees, and lateral bending right and left 30 degrees. Foraminal compression test and Spurling's test were positive. Reflexes were normal. A request was made for Temazepam 3mg, #30; Trazodone 50mg, #90; Valium 10mg, #120; Soma 350mg, #90; and Percocet 10/325mg, #120.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Temazepam 3 MG #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 24.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

Decision rationale: According to the MTUS, Benzodiazepines are not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks. Their range of action includes sedative/hypnotic, anxiolytic, anticonvulsant, and muscle relaxant. Chronic benzodiazepines are the treatment of choice in very few conditions. Tolerance to hypnotic effects develops rapidly. Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. A more appropriate treatment for anxiety disorder is an antidepressant. Tolerance to anticonvulsant and muscle relaxant effects occurs within weeks. (Baillargeon, 2003) (Ashton, 2005). According to the records, the injured worker has been taking his medication chronically. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Trazodone 50 MG #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Antidepressants Page(s): 15.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Trazodone: ODG, Mental Illness and Stress, Trazodone.

Decision rationale: Recommended as an option for insomnia, only for injured workers with potentially coexisting mild psychiatric symptoms such as depression or anxiety. See also Insomnia treatment, where it says there is limited evidence to support its use for insomnia, but it may be an option in injured workers with coexisting depression. Evidence for the off-label use of trazodone for treatment of insomnia is weak. The current recommendation is to utilize a combined pharmacologic and psychological and behavior treatment when primary insomnia is diagnosed. Also worth noting, there has been no dose-finding study performed to assess the dose of trazodone for insomnia in non-depressed injured workers. Other pharmacologic therapies should be recommended for primary insomnia before considering trazodone, especially if the insomnia is not accompanied by comorbid depression or recurrent treatment failure. There is no clear-cut evidence to recommend trazodone first line to treat primary insomnia. Therefore, at this time the requirements for treatment have not been met, and medical necessity has not been established.

Valium 10 MG #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 24.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

Decision rationale: According to the MTUS, Benzodiazepines are not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks. Their range of action includes sedative/hypnotic, anxiolytic, anticonvulsant, and muscle relaxant. Chronic benzodiazepines are the treatment of choice in very few conditions. Tolerance to hypnotic effects develops rapidly. Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. A more appropriate treatment for anxiety disorder is an antidepressant. Tolerance to anticonvulsant and muscle relaxant effects occurs within weeks. (Baillargeon, 2003) (Ashton, 2005). According to the records, the injured worker has been taking his medication chronically. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Soma 350 MG #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 65.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Soma Page(s): 29.

Decision rationale: According to the MTUS, Carisoprodol is not recommended. This medication is not indicated for long-term use. Carisoprodol is a commonly prescribed, centrally acting skeletal muscle relaxant whose primary active metabolite is meprobamate (a schedule-IV controlled substance). Carisoprodol is now scheduled in several states but not on a federal level. It has been suggested that the main effect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedative and relaxant effects. In regular abusers the main concern is the accumulation of meprobamate. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs. This includes the following: (1) increasing sedation of benzodiazepines or alcohol; (2) use to prevent side effects of cocaine; (3) use with tramadol to produce relaxation and euphoria; (4) as a combination with hydrocodone, an effect that some abusers claim is similar to heroin (referred to as a "Las Vegas Cocktail"); & (5) as a combination with codeine (referred to as "Soma Coma"). (Reeves, 1999) (Reeves, 2001) (Reeves, 2008) (Schears, 2004) There was a 300% increase in numbers of emergency room episodes related to carisoprodol from 1994 to 2005. (DHSS, 2005) Intoxication appears to include subdued consciousness, decreased cognitive function, and abnormalities of the eyes, vestibular function, appearance, gait and motor function. Intoxication includes the effects of both carisoprodol and meprobamate, both of which act on different neurotransmitters. (Bramness, 2007) (Bramness, 2004) A withdrawal syndrome has been documented that consists of insomnia, vomiting, tremors, muscle twitching, anxiety, and ataxia when abrupt discontinuation of large doses occurs. This is similar to withdrawal from meprobamate. (Reeves, 2007) (Reeves, 2004) There is little research in terms of weaning of high dose carisoprodol and there is no standard treatment regimen for injured workers with known dependence. Most treatment includes treatment for symptomatic complaints of withdrawal. Another option is to switch to phenobarbital to prevent withdrawal with subsequent tapering. A maximum dose of phenobarbital is 500 mg/day and the taper is 30 mg/day with a slower taper in an out injured worker setting. Tapering should be individualized for each injured worker. (Boothby, 2003) For more information and references, see Muscle relaxants. See also Weaning of medications. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Percocet 10/325 MG #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines
Page(s): 92.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines
Opioids Page(s): 74-97.

Decision rationale: According to the MTUS Chronic Pain Medical Treatment Guidelines section on Opioids, On-Going Management, p 74-97, (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the injured worker's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the injured worker's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain injured workers on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the injured worker should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or injured worker treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Additionally, the MTUS states that continued use of opioids requires (a) the injured worker has returned to work, (b) the injured worker has improved functioning and pain. There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.