

Case Number:	CM15-0169302		
Date Assigned:	09/16/2015	Date of Injury:	03/02/2005
Decision Date:	11/09/2015	UR Denial Date:	08/19/2015
Priority:	Standard	Application Received:	08/27/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, Indiana, Oregon
 Certification(s)/Specialty: Orthopedic Surgery

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 53 year old male who sustained an industrial injury on 03-02-2005. According to a report dated 07-23-2015, the injured worker was "doing poorly" and reported locking and catching of his left knee. The injured worker was in moderate distress. He had medial joint line tenderness about his left knee with grinding and catching. MRI scan of the left knee showed a large vertical split tear of the medial meniscus. Diagnosis included clinical and MRI scan evidence of a large vertical split tear of the medial meniscus of the left knee. Treatment to date has included physical therapy, medications, injections, bracing and rest. The injured worker remained disabled. The treatment plan included a request for authorization to proceed with a diagnostic and operative arthroscopy of the left knee with a partial medial meniscectomy. On 08-19-2015, Utilization review non-certified the request for assistant PA, preoperative lab: PT, PTT, preoperative urinalysis, preoperative EKG, preoperative chest x-ray and modified the request for twelve sessions of post-operative physical therapy and authorized the request for left knee arthroscopy with partial medial meniscectomy, preoperative medical clearance, cold therapy unit, Mobicrutches and preoperative labs: CBC and BMP.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Assistant PA: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Centers for Medicare and Medicaid services, Physician Fee Schedule Search, CPT code 29881.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) low back.

Decision rationale: CA MTUS/ACOEM is silent on the issue of a surgical assistant. ODG low back is referenced. More complex cases based off CPT code are felt to warrant the use of a surgical assistant. The requested procedure is knee arthroscopy. Given the level of complexity of the surgery, it is not medically necessary to have an assistant.

Preoperative lab: PT/PTT: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Institute for Clinical Systems Improvement (ICSI). Preoperative evaluation. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2006 Jul. 33 p. [37 references].

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) low back.

Decision rationale: CA MTUS/ACOEM is silent on the issue of preoperative clearance and testing. ODG, Low back, Preoperative testing general, is utilized. This chapter states that preoperative testing is guided by the patient's clinical history, comorbidities and physical examination findings. ODG states, "These investigations can be helpful to stratify risk, direct anesthetic choices, and guide postoperative management, but often are obtained because of protocol rather than medical necessity. The decision to order preoperative tests should be guided by the patient's clinical history, comorbidities and physical examination findings. Patients with signs or symptoms of active cardiovascular disease should be evaluated with appropriate testing, regardless of their preoperative status." Preoperative ECG in patients without known risk factor for coronary artery disease, regardless of age, may not be necessary. CBC is recommended for surgeries with large anticipated blood loss. Creatinine is recommended for patient with renal failure. Electrocardiography is recommended for patients undergoing high-risk surgery and those undergoing intermediate risk surgery who have additional risk factors. Patients undergoing low risk surgery do not require electrocardiography. Based on the information provided for review, there is no indication of any of these clinical scenarios present in this case. In this case the patient is healthy without comorbidities or physical examination findings concerning to warrant preoperative testing prior to the proposed surgical procedure. Therefore, the request is not medically necessary.

Preoperative urinalysis: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Bibliographic Source: National Collaborating

Centre for Acute Care. Preoperative tests: the use of routine preoperative tests for elective surgery: evidence, methods & guidance. London (UK).

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) low back.

Decision rationale: CA MTUS/ACOEM is silent on the issue of preoperative clearance and testing. ODG, Low back, Preoperative testing general, is utilized. This chapter states that preoperative testing is guided by the patient's clinical history, comorbidities and physical examination findings. ODG states, "These investigations can be helpful to stratify risk, direct anesthetic choices, and guide postoperative management, but often are obtained because of protocol rather than medical necessity. The decision to order preoperative tests should be guided by the patient's clinical history, comorbidities and physical examination findings. Patients with signs or symptoms of active cardiovascular disease should be evaluated with appropriate testing, regardless of their preoperative status." Preoperative ECG in patients without known risk factor for coronary artery disease, regardless of age, may not be necessary. CBC is recommended for surgeries with large anticipated blood loss. Creatinine is recommended for patient with renal failure. Electrocardiography is recommended for patients undergoing high-risk surgery and those undergoing intermediate risk surgery who have additional risk factors. Patients undergoing low risk surgery do not require electrocardiography. Based on the information provided for review, there is no indication of any of these clinical scenarios present in this case. In this case the patient is healthy without comorbidities or physical examination findings concerning to warrant preoperative testing prior to the proposed surgical procedure. Therefore, the request is not medically necessary.

Preoperative EKG: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Institute for Clinical Systems Improvement (ICSI). Preoperative evaluation. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2006 Jul. 33 p. [37 references].

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) low back.

Decision rationale: CA MTUS/ACOEM is silent on the issue of preoperative clearance and testing. ODG, Low back, Preoperative testing general, is utilized. This chapter states that preoperative testing is guided by the patient's clinical history, comorbidities and physical examination findings. ODG states these investigations can be helpful to stratify risk, direct anesthetic choices, and guide postoperative management, but often are obtained because of protocol rather than medical necessity. The decision to order preoperative tests should be guided by the patient's clinical history, comorbidities and physical examination findings. Patients with signs or symptoms of active cardiovascular disease should be evaluated with appropriate testing, regardless of their preoperative status. Preoperative ECG in patients without known risk factor for coronary artery disease, regardless of age, may not be necessary. CBC is recommended for surgeries with large anticipated blood loss. Creatinine is recommended for patient with renal

failure. Electrocardiography is recommended for patients undergoing high-risk surgery and those undergoing intermediate risk surgery who have additional risk factors. Patients undergoing low risk surgery do not require electrocardiography. Based on the information provided for review, there is no indication of any of these clinical scenarios present in this case. In this case the patient is healthy without comorbidities or physical examination findings concerning to warrant preoperative testing prior to the proposed surgical procedure. Therefore, the request is not medically necessary.

Preoperative chest x-ray: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Institute for Clinical Systems Improvement (ICSI). Preoperative evaluation. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2006 Jul. 33 p. [37 references].

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) low back.

Decision rationale: CA MTUS/ACOEM is silent on the issue of preoperative clearance and testing. ODG, Low back, Preoperative testing general, is utilized. This chapter states that preoperative testing is guided by the patient's clinical history, comorbidities and physical examination findings. ODG states these investigations can be helpful to stratify risk, direct anesthetic choices, and guide postoperative management, but often are obtained because of protocol rather than medical necessity. The decision to order preoperative tests should be guided by the patient's clinical history, comorbidities and physical examination findings. Patients with signs or symptoms of active cardiovascular disease should be evaluated with appropriate testing, regardless of their preoperative status. Preoperative ECG in patients without known risk factor for coronary artery disease, regardless of age, may not be necessary. CBC is recommended for surgeries with large anticipated blood loss. Creatinine is recommended for patient with renal failure. Electrocardiography is recommended for patients undergoing high-risk surgery and those undergoing intermediate risk surgery who have additional risk factors. Patients undergoing low risk surgery do not require electrocardiography. Based on the information provided for review, there is no indication of any of these clinical scenarios present in this case. In this case the patient is healthy without comorbidities or physical examination findings concerning to warrant preoperative testing prior to the proposed surgical procedure. Therefore, the request is not medically necessary.

Twelve sessions of post operative physical therapy: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Postsurgical Treatment 2009, Section(s): Knee.

Decision rationale: According to the CA MTUS/Post Surgical Treatment Guidelines, Knee Meniscectomy, page 24, 12 visits of therapy are recommended after arthroscopy with partial meniscectomy over a 12-week period. The guidelines recommend initially of the 12 visits to be performed. As the request exceeds the initial allowable visits, the request is not medically necessary.