

Case Number:	CM15-0170554		
Date Assigned:	09/11/2015	Date of Injury:	10/19/1995
Decision Date:	10/13/2015	UR Denial Date:	07/30/2015
Priority:	Standard	Application Received:	08/31/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 61 year old male with an industrial injury dated 10-19-1995. Medical record review indicates he is being treated for lumbar degenerative disc disease, discogenic low back pain, chronic lumbar 5-sacral 1 radiculitis, chronic low back pain and lumbar myofascial pain. He presented on 07-21-2015 for reevaluation of his low back pain with burning and numbness in the lateral and anterior thighs. Pain levels are documented as 8-9 out of 10 before medication, coming down to 5-6 with medication. Documentation also notes with medication he is active, rides a bike and does house and yard work. His medications included Percocet, Diclofenac and Omeprazole (for gastrointestinal upset from his medications.) Physical exam of the lumbar spine noted range of motion is full flexion and extension was decreased to 15 degrees causing pain. Work status is documented as retired. The progress note dated 04-28-2015 also documented low back pain with pain levels as 7-9 out of 10 without medication coming down to 4-6 out of 10 with medication. Medications are documented as Percocet and Prilosec. The provider documented the injured worker "takes a rare Percocet." "The last time he was given a prescription was January 2015." Documentation notes a signed opioid agreement in the chart and CURES report from 07-08-2015 was consistent. Prior treatments are documented as lumbar epidural steroid injections and medications. The treatment request is for Prilosec 20 mg #30 with 4 refills and Percocet 5-325 mg #30. The request for Prilosec 20 mg # 30 with 4 refills was non- certified by utilization review. The request for Percocet 5-325 mg was modified to 15 tablets of Percocet 5-325 mg by utilization review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Percocet 5/325mg #30: Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids for chronic pain.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work; (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. These criteria are met in the proved documentation for review and the request is therefore medically necessary.

Prilosec 20mg #30 with 4 refills: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy and proton pump inhibitors (PPI) states: Recommend with precautions as indicated below. Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or a anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastro duodenal lesions. Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g., ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg Omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. There is no documentation provided that places this patient at intermediate or high risk that would justify the use of a PPI. There is no mention of current gastrointestinal or cardiovascular disease. For these reasons, the criteria set forth above per the California MTUS for the use of this medication has not been met. Therefore, the request is not medically necessary.