

Case Number:	CM15-0171654		
Date Assigned:	09/11/2015	Date of Injury:	06/03/2003
Decision Date:	11/25/2015	UR Denial Date:	08/18/2015
Priority:	Standard	Application Received:	08/31/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Emergency Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 63-year-old male, who sustained an industrial injury on June 3, 2003, incurring upper back injuries. He was diagnosed with cervical degenerative disc disease, cervical radiculopathy, cervicgia and headaches. Treatments included physical therapy and home exercise program, pain medications, anti-inflammatory drugs, neuropathic medications, medial branch block, and work restrictions. He underwent a cervical laminectomy and cervical fusion on January 3, 2004. Currently, the injured worker complained of persistent neck pain, tenderness and muscle spasms. The injured worker had difficulty with sleeping secondary to the chronic pain. He rated his neck pain with medications 5 out of 10 on a pain scale of 1 to 10 and without medications 8 out of 10. He noted that the medial branch block provided 100% pain relief. The injured worker noted that his prescribed medications help with his functional mobility. On April 27, 2015, a cervical Magnetic Resonance Imaging revealed severe right neural foraminal stenosis. His cervical pain limited his activities of daily living. The treatment plan that was requested for authorization on August 25, 2015, included prescriptions for Vicodin 5-300 mg, #30; Ultram 50 mg, #90; Gabapentin 300 mg, #90; and Relafen 750 mg, #60. On August 18, 2015, a request for prescriptions for Vicodin, Ultram, Gabapentin and Relafen was denied by utilization review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Vicodin 5/300mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, criteria for use.

Decision rationale: The request is for the use of a medication in the opioid class. The MTUS guidelines state that for ongoing treatment with a pharmaceutical in this class, certain requirements are necessary. This includes not only adequate pain control, but also functional improvement. Four domains have been proposed for management of patients on opioids. This includes pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant drug-related behaviors. In this case, there is inadequate documentation of persistent functional improvement seen. As such, the request is not certified. All opioid medications should be titrated down slowly in order to prevent a significant withdrawal syndrome. Therefore, the requested treatment is not medically necessary.

Ultram 50mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids for chronic pain.

Decision rationale: Tramadol is a pain medication in the category of a centrally acting analgesic. They exhibit opioid activity and a mechanism of action that inhibits the reuptake of serotonin and norepinephrine. Centrally acting drugs are reported to be effective in managing neuropathic type pain although it is not recommended as first line therapy. The side-effect profile is similar to opioids. For chronic back pain, it appears to be efficacious for short-term pain relief, but long term (>16 weeks) results are limited. It also did not appear to improve function. The use of Tramadol for osteoarthritis is indicated for short-term use only (<3 months) with poor long-term benefit. In this case, the patient does not meet the qualifying criteria. This is secondary to the duration of use, with this medication being indicated on a short-term basis only. As such, the request is not certified. Therefore, the requested treatment is not medically necessary.

Gabapentin 300mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Antiepilepsy drugs (AEDs).

Decision rationale: The request is for the use of a medication in the category of an anti-epileptic drug (AED). These medications are recommended for certain types of neuropathic pain. Most of the randomized clinical control trials involved include post-herpetic neuralgia and painful polyneuropathy such as in diabetes. There are few trials, which have studied central pain or radiculopathy. The MTUS guidelines state that a good response to treatment is 50% reduction in pain. At least a 30% reduction in pain is required for ongoing use, and if this is not seen, this should trigger a change in therapy. There also should be documentation of functional

improvement and side effects incurred with use. Disease states, which prompt use of these medications, include post-herpetic neuralgia, spinal cord injury, chronic regional pain syndrome, lumbar spinal stenosis, post-operative pain, and central pain. There is inadequate evidence to support use in non-specific axial low back pain or myofascial pain. In this case, there is documentation of neuropathic pain. In addition to pain control, functional improvement is necessary for continued use. The records also do not reveal functional improvement. As such, the request is not certified. Therefore, the requested treatment is not medically necessary.

Relafen 750mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steroidal anti-inflammatory drugs).

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic)/NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: The request is for the use of a medication in the NSAID class. The ODG state the following regarding this topic: Specific recommendations: Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxen being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain - Acute low back pain & acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting to negative evidence that NSAIDs are more effective than acetaminophen for acute LBP. (van Tulder, 2006) (Hancock, 2007) For patients with acute low back pain with sciatica a recent Cochrane review (including three heterogeneous randomized controlled trials) found no differences in treatment with NSAIDs vs. placebo. In patients with axial low back pain, this same review found that NSAIDs were not more effective than acetaminophen for acute low-back pain, and that acetaminophen had fewer side effects. (Roelofs-Cochrane, 2008) The addition of NSAIDs or spinal manipulative therapy does not appear to increase recovery in patients with acute low back pain over that received with acetaminophen treatment and advice from their physician. (Hancock, 2007) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough pain and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in patients with neuropathic pain. (Namaka, 2004) (Gore, 2006) See

NSAIDs, GI symptoms & cardiovascular risk; NSAIDs, hypertension and renal function; & Medications for acute pain (analgesics). Besides the above well-documented side effects of NSAIDs, there are other less well-known effects of NSAIDs, and the use of NSAIDs has been shown to possibly delay and hamper healing in all the soft tissues, including muscles, ligaments, tendons, and cartilage. (Maroon, 2006) The risks of NSAIDs in older patients, which include increased cardiovascular risk and gastrointestinal toxicity, may outweigh the benefits of these medications. (AGS, 2009) As stated above, acetaminophen would be considered first-line treatment for chronic pain. In this case, the continued use of an NSAID is not supported. This is secondary to inadequate documentation of functional improvement benefit seen. In addition, the duration of use places the patient at risk for gastrointestinal and cardiovascular side-effects. In addition, it is known that use of NSAIDs delays the healing of soft tissue including ligaments, tendons, and cartilage. As such, the request is not medically necessary.