MAXIMUS Federal Services

Case Number:	CM15-0172708		
Date Assigned:	09/14/2015	Date of Injury:	08/25/2014
Decision Date:	10/14/2015	UR Denial Date:	07/28/2015
Priority:	Standard	Application Received:	09/01/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: North Carolina
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 27 year old male, who sustained an industrial injury on 8-25-2014. The medical records indicate that the injured worker is undergoing treatment for left elbow sprain-strain, left lateral epicondylitis, left forearm strain, left forearm muscle spasm, left wrist tenosynovitis, status post-surgery left wrist. According to the progress report dated 7-21-2015, the injured worker complains of frequent, mild throbbing left elbow pain, frequent, moderate achy left forearm pain, and frequent, mild achy left wrist pain. The level of pain is not rated. The physical examination of the left elbow reveals tenderness to palpation of the anterior elbow, lateral elbow, medial elbow, and posterior elbow. There is muscle spasm of the dorsal and volar forearm. Examination of the left forearm reveals tenderness to palpation of the dorsal forearm, radial forearm, ulnar forearm, and volar forearm. Range of motion is limited with pronation (33 degrees). Examination of the left wrist reveals tenderness to palpation over the volar wrist. The current medications are Norco. There is documentation of ongoing treatment with Norco since at least 2-19-2015. Treatment to date has included medication management, x-rays, electrodiagnostic testing, injection therapy, and surgical intervention. Work status is described as off work. The original utilization review (7-28-2015) partially approved a request for Norco #45 (original request for #60). The request for trigger point impedance imaging and unknown localized intense neurostimulation therapy was non-certified.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids for chronic pain.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain dairy that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work; (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant improvement in VAS scores for significant periods of time. There are no objective measurements of improvement in function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

One trigger points impedance imaging: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Trigger point injections.

Decision rationale: The California chronic pain medical treatment guidelines section on trigger point injections states: Trigger point injections: Recommended only for myofascial pain syndrome as indicated below, with limited lasting value. Not recommended for radicular pain. Trigger point injections with an anesthetic such as bupivacaine are recommended for non-resolving trigger points, but the addition of a corticosteroid is not generally recommended. Not recommended for radicular pain. A trigger point is a discrete focal tenderness located in a palpable taut band of skeletal muscle, which produces a local twitch in response to stimulus to the band. Trigger points may be present in up to 33-50% of the adult population. Myofascial pain syndrome is a regional painful muscle condition with a direct relationship between a specific trigger point and its associated pain region. These injections may occasionally be necessary to maintain function in those with myofascial problems when myofascial trigger points are present on examination. Not recommended for typical back pain or neck pain. (Graff-Radford, 2004) (Nelemans-Cochrane, 2002) For fibromyalgia syndrome, trigger point injections have not been proven effective. (Goldenberg, 2004) Criteria for the use of Trigger point injections: Trigger point injections with a local anesthetic may be recommended for the treatment of chronic low back or neck pain with myofascial pain syndrome when all of the following criteria are met:(1) Documentation of circumscribed trigger points with evidence upon palpation of a twitch response as well as referred pain; (2) Symptoms have persisted for more than three months; (3) Medical management therapies such as ongoing stretching exercises, physical therapy, NSAIDs and muscle relaxants have failed to control pain; (4) Radiculopathy is not present (by exam, imaging, or neuro-testing); (5) Not more than 3-4 injections per session; (6) No repeat injections unless a greater than 50% pain relief is obtained for six weeks after an injection and there is documented evidence of functional improvement; (7) Frequency should not be at an interval less than two months; (8) Trigger point injections with any substance (e.g., saline or glucose) other than local anesthetic with or without steroid are not recommended. (Colorado, 2002) (BlueCross BlueShield, 2004) The provided clinical documentation fails to show circumscribed trigger points with evidence upon palpation of a twitch response as well as referred pain. Therefore, criteria have not been met and the request is not certified as if injection is not medically necessary, neither would imaging.

Unknown localized intense neurostimulation therapy: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Transcutaneous electrotherapy.

Decision rationale: The California MTUS section on NMES states: Neuromuscular electrical stimulation (NMES devices) not recommended. NMES is used primarily as part of a rehabilitation program following stroke and there is no evidence to support its use in chronic pain. There are no intervention trials suggesting benefit from NMES for chronic pain. (Moore, 1997) (Gaines, 2004) The scientific evidence related to electromyography (EMG)-triggered electrical stimulation therapy continues to evolve, and this therapy appears to be useful in a supervised physical therapy setting to rehabilitate atrophied upper extremity muscles following stroke and as part of a comprehensive PT program. Neuromuscular Electrical Stimulation Devices (NMES), NMES, through multiple channels, attempts to stimulate motor nerves and alternately causes contraction and relaxation of muscles, unlike a TENS device which is intended to alter the perception of pain. NMES devices are used to prevent or retard disuse atrophy, relax muscle spasm, increase blood circulation, maintain or increase range-of-motion, and re-educate muscles. Functional neuromuscular stimulation (also called electrical neuromuscular stimulation and EMG-triggered neuromuscular stimulation) attempts to replace stimuli from destroyed nerve pathways with computer-controlled sequential electrical stimulation of muscles to enable spinal cord-injured or stroke patients to function independently, or at least maintain healthy muscle tone and strength. Also used to stimulate quadriceps muscles following major knee surgeries to maintain and enhance strength during rehabilitation. (BlueCross BlueShield, 2005) (Aetna, 2005) The patient does not have the diagnosis of stroke and therefore the request is not medically necessary.