MAXIMUS Federal Services

Case Number:	CM15-0173062		
Date Assigned:	09/15/2015	Date of Injury:	04/22/2002
Decision Date:	10/14/2015	UR Denial Date:	08/20/2015
Priority:	Standard	Application Received:	09/02/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: North Carolina
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 61-year-old female who sustained an industrial injury April 22, 2002. Diagnoses have included low back pain and sciatica. Documented treatment includes 10 visits of Acupuncture with the last being February 25, 2015. No other treatments are present in provided documentation. The treating physician's plan of care includes 6 sessions of acupuncture for the low back and Relafen, which was denied August 20, 2015. Current work status is not available.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Acupuncture for low back qty: 6: Upheld

Claims Administrator guideline: Decision based on MTUS Acupuncture Treatment 2007.

MAXIMUS guideline: Decision based on MTUS Acupuncture Treatment 2007.

Decision rationale: The California chronic pain medical treatment guidelines section on acupuncture states: Acupuncture is used as an option when pain medication is reduced or not tolerated, it may be used as an adjunct to physical rehabilitation and/or surgical intervention to hasten functional recovery. It is the insertion and removal of filiform needles to stimulate

acupoints (acupuncture points). Needles may be inserted, manipulated, and retained for a period of time. Acupuncture can be used to reduce pain, reduce inflammation, increase blood flow, increase range of motion, decrease the side effect of medication-induced nausea, promote relaxation in an anxious patient, and reduce muscle spasm. Time to produce functional improvement is 3-6 treatments and frequency is 1-3 times per week. The requested amount of session is in excess of the recommendation unless improvement is noted by 3-6 sessions. As previous session have not produced documented objective improvement in pain and function. Therefore, the request is not medically necessary.

Relafen: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steriodal anti-inflammatory drugs).

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy states: Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. This medication is recommended for the shortest period of time and at the lowest dose possible. The definition of shortest period possible is not clearly defined in the California MTUS. However, there is no quantity or dosing information given in the request and therefore it is not medically necessary.