

Case Number:	CM15-0173813		
Date Assigned:	09/15/2015	Date of Injury:	08/14/2014
Decision Date:	10/21/2015	UR Denial Date:	08/21/2015
Priority:	Standard	Application Received:	09/03/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, District of Columbia, Maryland
 Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 73-year-old male who sustained an industrial injury August 14, 2014. Diagnoses have included shoulder sprain, and MRI of May 15, 2015 showed complete tear of supraspinatus tendon with 45 mm tendinous retraction and acromioclavicular osteoarthritis of the left shoulder. Documented treatment includes an unspecified number of physical therapy and acupuncture sessions, home exercise, group therapy, biofeedback therapy, and medication, but the injured worker continues to present with "constant" neck and left shoulder pain radiating down his arm. He reports weakness, numbness to the upper extremity, and pain becomes worse when reaching with reported rating of 7 out of 10. August 14, 2015 examination revealed left abduction and forward flexion 125 degrees; internal rotation and extension 65 degrees, and adduction and back extension 10 degrees. Right shoulder range was stated to be "within normal limits." The treating physician's plan of care includes compound creams: Flurbiprofen 25 percent, Cyclobenzaprine 2 percent; and, Gabapentin 15 percent, Dextromethorphan 10 percent, Amitriptyline 4 percent, which was denied August 21, 2015. He has not been working.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Flurbiprofen 25%, Cyclobenzaprine 2% compound cream 180gms: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: Per MTUS with regard to Flurbiprofen (p112), "These medications may be useful for chronic musculoskeletal pain, but there are no long-term studies of their effectiveness or safety. (Mason, 2004) Indications: Osteoarthritis and tendinitis, in particular, that of the knee and elbow or other joints that are amenable to topical treatment: Recommended for short-term use (4-12 weeks). There is little evidence to utilize topical NSAIDs for treatment of osteoarthritis of the spine, hip or shoulder." Flurbiprofen may be indicated. Per MTUS CPMTG p113, "There is no evidence for use of any other muscle relaxant as a topical product. [besides baclofen, which is also not recommended]" Cyclobenzaprine is not indicated. Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. As topical cyclobenzaprine is not recommended, the compound is not medically necessary.

Gabapentin 15%, Dextromethorphan 10%, Amitriptyline 4% compound cream 180gms:
Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: Per MTUS p113 with regard to topical gabapentin: "Not recommended. There is no peer-reviewed literature to support use." Per the article "Topical Analgesics in the Management of Acute and Chronic Pain" published in Mayo Clinic Proceedings (Vol 88, Issue 2, p 195-205), "Studies in healthy volunteers demonstrated that topical amitriptyline at concentrations of 50 and 100 mmol/L produced a significant analgesic effect ($P < .05$) when compared with placebo and was associated with transient increases in tactile and mechanical nociceptive thresholds." Amitriptyline may be indicated. The CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of dextromethorphan. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status

equivalent to "not recommended". Since dextromethorphan and gabapentin are not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually. The request is not medically necessary.