

Case Number:	CM15-0175041		
Date Assigned:	09/16/2015	Date of Injury:	06/20/2008
Decision Date:	10/19/2015	UR Denial Date:	08/18/2015
Priority:	Standard	Application Received:	09/04/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The 61-year-old male injured worker suffered an industrial injury on 6-20-2008. The diagnoses included radiculopathy. On 8-10-2015 the treating provider reported back pain radiating from the low back down both legs rated with medication as 4 out of 10 and without medication 6 out of 10. The quality of sleep was poor. There was a permanent spinal cord stimulator implantation on 7-21-2015 and had been able to decrease the Opana 30mg from 3 per day to 2 per day and decreased Norco from 6 per day to 4 per day. He stated that it decreased the pain by greater than 50%. On exam, there was an altered gait. The lumbar spine had restricted range of motion and positive lumbar facet loading signs. The provider noted the injured worker had difficulty translating pain gains into increased activity at least partially related to ongoing depression and fear avoidance and deconditioning after 7 years of inactivity make him appropriate for a functional restoration program. Prior treatments were injections, opiates, and extensive physical therapy. The Utilization Review on 8-18-2015 determined non-certification/modification for Consultation for possible multidisciplinary pain treatment program, Opana ER 30mg qty: 90 and Tizanidine HCl 2mg qty: 90.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Consultation for possible multidisciplinary pain treatment program: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Functional restoration programs (FRPs).

Decision rationale: The California chronic pain medical treatment guidelines section on functional restoration programs states: Recommended, although research is still ongoing as to how to most appropriately screen for inclusion in these programs. Functional restoration programs (FRPs), a type of treatment included in the category of interdisciplinary pain programs (see Chronic pain programs), were originally developed by Mayer and Gatchel. FRPs were designed to use a medically directed, interdisciplinary pain management approach geared specifically to patients with chronic disabling occupational musculoskeletal disorders. These programs emphasize the importance of function over the elimination of pain. FRPs incorporate components of exercise progression with disability management and psychosocial intervention. Long-term evidence suggests that the benefit of these programs diminishes over time, but remains positive when compared to cohorts that did not receive an intensive program. (Bendix, 1998) A Cochrane review suggests that there is strong evidence that intensive multidisciplinary rehabilitation with functional restoration reduces pain and improves function of patients with low back pain. The evidence is contradictory when evaluating the programs in terms of vocational outcomes. (Guzman 2001) It must be noted that all studies used for the Cochrane review excluded individuals with extensive radiculopathy, and several of the studies excluded patients who were receiving a pension, limiting the generalizability of the above results. Studies published after the Cochrane review also indicate that intensive programs show greater effectiveness, in particular in terms of return to work, than less intensive treatment. (Airaksinen, 2006) There appears to be little scientific evidence for the effectiveness of multidisciplinary biopsychosocial rehabilitation compared with other rehabilitation facilities for neck, shoulder pain, as opposed to low back pain, and generalized pain syndromes. (Karjalainen, 2003) Treatment is not suggested for longer than 2 weeks without evidence of demonstrated efficacy as documented by subjective and objective gains. For general information, see Chronic pain programs. While functional restoration programs are recommended per the California MTUS, the length of time is for 2 weeks unless there is documentation of demonstrated efficacy by subjective and objective gains. The request is for a single evaluation and therefore is medically necessary.

Opana ER 30mg qty: 90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids for chronic pain.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a)

Prescriptions from a single practitioner taken as directed, and all prescriptions from single pharmacy.(b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the 4 A's (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids(a) If the patient has returned to work(b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant improvement in VAS scores for significant periods of time with pain only decreasing from a 6/10 to a 4/10 with medications. There are no objective measurements of improvement in function or activity specifically due to the medication. The patient is not working. Therefore, not all criteria for the ongoing use of opioids have been met and the request is not medically necessary.

Tizanidine HCl 2mg qty: 90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

Decision rationale: The California chronic pain medical treatment guidelines section on muscle relaxants states: Recommend non-sedating muscle relaxants with caution as a second-line option

for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. In addition, there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) (Chou, 2004) This medication is not intended for long-term use per the California MTUS. The medication has not been prescribed for the flare-up of chronic low back pain, but rather for ongoing and chronic lumbar pain. This is not an approved use for the medication. For these reasons, criteria for the use of this medication have not been met. Therefore, the request is not medically necessary.