

<b>Case Number:</b>	CM15-0175323		
<b>Date Assigned:</b>	09/25/2015	<b>Date of Injury:</b>	11/10/2009
<b>Decision Date:</b>	11/19/2015	<b>UR Denial Date:</b>	08/05/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	09/04/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: California  
 Certification(s)/Specialty: Emergency Medicine

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51 year old female, who sustained an industrial injury on 1-10-2009. Medical records indicate the worker is undergoing treatment for lumbosacral radiculopathy, insomnia, cervical radiculitis, left shoulder impingement and status post left carpal tunnel release. A recent progress report dated 7-8-2015, reported the injured worker complained of increased lumbosacral pain with radiation to bilateral lower extremities. The same complaints have been consistent since at least 3-4-2015. Physical examination revealed "decreased lumbar range of motion with positive straight leg raise". Treatment to date has included medications that include Cyclobenzaprine, Hydrocodone-Acetaminophen, Hydrocodone-APAP and Lorzone. On 7-21-2015, the Request for Authorization requested Cyclobenzaprine 10mg #60, Hydrocodone- APAP 10-325mg #60 (since at least 3-4-2015), Lorzone 750mg #60 and Hydrocodone- acetaminophen 10mg #90. On 8-5-2015, the Utilization Review noncertified the request for Cyclobenzaprine 10mg #60, Hydrocodone-APAP 10-325mg #60 and Lorzone 750mg #60 and modified the request for Hydrocodone-acetaminophen 10mg #90 to #60.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Cyclobenzaprine 10mg #60:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

**Decision rationale:** The request is for the use of a muscle relaxant to aid in pain relief. The MTUS guidelines state that the use of a medication in this class is indicated as a second-line option for short-term treatment of acute exacerbations of low back pain. Muscle relaxants may be effective in reducing pain and muscle tension, which can increase mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain improvement. Efficacy appears to diminish over time, and prolonged use may lead to dependence. (Homik, 2004) Due to inadequate documentation of a recent acute exacerbation and poor effectiveness for chronic long-term use, the request is not medically necessary.

**Hydrocodone-acetaminophen 10mg #90:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, criteria for use.

**Decision rationale:** The request is for the use of a medication in the opioid class. The MTUS guidelines state that for ongoing treatment with a pharmaceutical in this class, certain requirements are necessary. This includes not only adequate pain control, but also functional improvement. Four domains have been proposed for management of patients on opioids. This includes pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant drug-related behaviors. In this case, there is inadequate documentation of persistent functional improvement seen. As such, the request is not medically necessary. All opioid medications should be titrated down slowly in order to prevent a significant withdrawal syndrome.

**Hydrocodone-APAP 10/325mg #60:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, criteria for use.

**Decision rationale:** The request is for the use of a medication in the opioid class. The MTUS guidelines state that for ongoing treatment with a pharmaceutical in this class, certain requirements are necessary. This includes not only adequate pain control, but also functional improvement. Four domains have been proposed for management of patients on opioids. This includes pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant drug-related behaviors. In this case, there is inadequate documentation of persistent functional improvement seen. As such, the request is not medically necessary. All opioid medications should be titrated down slowly in order to prevent a significant withdrawal syndrome.

**Lorzone 750mg #60:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not cite any medical evidence for its decision.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

**Decision rationale:** The request is for the use of a muscle relaxant to aid in pain relief. The MTUS guidelines state that the use of a medication in this class is indicated as a second-line option for short-term treatment of acute exacerbations of low back pain. Muscle relaxants may be effective in reducing pain and muscle tension, which can increase mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain improvement. Efficacy appears to diminish over time, and prolonged use may lead to dependence. (Homik, 2004) Due to inadequate documentation of a recent acute exacerbation and poor effectiveness for chronic long-term use, the request is not medically necessary.