

Case Number:	CM15-0176049		
Date Assigned:	09/17/2015	Date of Injury:	06/09/2010
Decision Date:	10/19/2015	UR Denial Date:	08/13/2015
Priority:	Standard	Application Received:	09/08/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 40 year old female, who sustained an industrial injury on 6-09-2010, resulting from a slip and fall. The injured worker was diagnosed as having chronic cervical pain with cervical disc protrusion at C6-7 without spinal stenosis or neuroforaminal narrowing, chronic left shoulder pain, chronic thoracic myofascial pain, and chronic lumbosacral pain. Non-industrial conditions included fibromyalgia (currently not active), bipolar disorder, schizoaffective disorder, and depression (currently in remission). Treatment to date has included diagnostics, chiropractic, acupuncture, physical therapy, and medications. Qualified Medical Examination (6-17-2014) noted the use of Norco, 180 per month. Currently (7-07-2015), the injured worker complains of pain in her neck and upper back, and slight pain in her lower back. She denied having left shoulder pain or headaches at the time. Objective findings included paracervical tenderness from C2 to C7-T1. There was parathoracic tenderness from T1 to T10 and paralumbar tenderness from L2 to L5-S1. It was documented that she "gets pain relief from the Norco and improved functioning". She was documented as not having significant side effects from the medication and had "increased physical and psychosocial functioning" as a result of taking opiate medication. The Pain Disability Index questionnaire noted ratings of 4 with medication use and 7-8 without (compared to 3-5 with medication use and 6-8 without on 5-12-2015). There was no evidence of non-compliance with medication and a signed pain management agreement was on file. She was also taking Baclofen for muscle spasm (as provided by primary care doctor). Her work status remained modified. Urine toxicology was not submitted. The treatment plan included the use of Norco 5-325mg #180 (1 tab every 4-6 hours consistent for at least 6 months), modified to Norco 5-325mg #15 by Utilization Review on 8-13-2015.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 5/325 MG #180: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids for chronic pain.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004). The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is documented significant improvement in VAS scores for significant periods of time with pain decreasing from a 8/10 to a 4/10. There are no objective measurements of improvement in function or activity specifically due to the medication. Therefore, all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.