

<b>Case Number:</b>	CM15-0177799		
<b>Date Assigned:</b>	11/03/2015	<b>Date of Injury:</b>	06/30/2014
<b>Decision Date:</b>	12/14/2015	<b>UR Denial Date:</b>	08/11/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	09/09/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following  
 credentials: State(s) of Licensure: North Carolina  
 Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 33 year old female, who sustained an industrial injury on 6-30-14. The injured worker is diagnosed with lumbar facet joint arthropathy, sacroiliac joint inflammation, myofascial pain, lumbar intervertebral disc degeneration and chronic pain. Her work status is full duty. Notes dated 7-14-15 and 7-24-15 reveals the injured worker presented with complaints of intermittent bilateral low back pain that radiates to the back of her thigh described as aching, electrical and sharp. She reports right leg weakness and numbness. The pain is increased with low back extension and flexion as well as walking and relieved by lying down and walking. Physical examinations dated 7-14-15 and 7-24-15 revealed tenderness to palpation over the lumbar paraspinal muscles and the facet joint on the right, as well as 2+ muscle spasms noted over the lower paraspinal muscles. Limited lumbar flexion and extension is noted. Treatment to date has included medications-Ibuprofen (1-2015), physical therapy note dated 7-28-15 states the injured worker is exhibiting "improved ability to isolate and engage "TrA" with cueing and coaching, no increased complaints of back pain with there-ex, continue to address limitation with range of motion, mobility, functional and core strength to aid in activities of daily living tolerance". Diagnostic studies include urine toxicology screen, lumbar x-rays, which revealed no evidence of instability, disc spaces are well preserved and no lytic or destructive lesion identified and a lumbar MRI, which revealed disc desiccation with annular bulging and small left posterior lateral disc protrusion with associated small annular fissure and mild facet arthropathy, per physician note dated 7-24-15. A request for authorization dated 7-28-15 for Ibuprofen 600 mg

#90 with 2 refills is modified to #90 with no refills and 6 physical therapy sessions is modified to 5 sessions, per Utilization Review letter dated 8-11-15.

### **IMR ISSUES, DECISIONS AND RATIONALES**

The Final Determination was based on decisions for the disputed items/services set forth below:

**Ibuprofen 600 mg #90 with 2 refills:** Overturned

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steriodal anti-inflammatory drugs).

**Decision rationale:** The California chronic pain medical treatment guidelines section on NSAID therapy states: Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. This medication is recommended for the shortest period of time and at the lowest dose possible. The dosing of this medication is within the California MTUS guideline recommendations. The definition of shortest period possible is not clearly defined in the California MTUS. Therefore the request is medically necessary.

**6 physical therapy sessions:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Physical Medicine.

**Decision rationale:** The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The patient has already completed physical therapy. The request is in excess of these recommendations per the California MTUS. There is no objective reason why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions in the provided clinical documentation. Therefore the request is not medically necessary.