

Case Number:	CM15-0178843		
Date Assigned:	09/21/2015	Date of Injury:	12/04/2009
Decision Date:	10/23/2015	UR Denial Date:	09/01/2015
Priority:	Standard	Application Received:	09/11/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 53 year old male, who sustained an industrial injury on 12-04-2009. The injured worker is being treated for lumbar disc injury, L5-S1 radiculopathy and bilateral sciatica. Treatment to date has included medications and diagnostics. Per the Follow-up Evaluation dated 8-19-2015 the injured worker reported back pain rated as 5-8 out of 10 in severity and pain with medication as 4-5 out of 10 in severity. Objective findings of the lumbar spine included left more than right L4-5 and L5-S1 moderate tenderness. Range of motion is complete in all directions with moderate pain upon extension, right lateral flexion with slight pain upon forward flexion and left lateral flexion. Per the medical records dated 3-31-2015 to 8-19-2015 there is no documentation of improvement in symptoms, increase in activities of daily living or decrease in pain level with the current treatment. Work status was not provided on this date. The plan of care included, and authorization was requested for Lidoderm patch 5% #90 and 6 psychologist sessions. On 9-01-2015, Utilization Review non-certified the request for Lidoderm patch 5% #90 and 6 psychologist sessions citing lack of established medical necessity.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Lidoderm patch 5% #90 with 6 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The California chronic pain medical treatment guidelines section on topical lidocaine states: Lidocaine Indication: Neuropathic pain; Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Topical lidocaine, in the formulation of a dermal patch (Lidoderm) has been designated for orphan status by the FDA for neuropathic pain. Lidoderm is also used off-label for diabetic neuropathy. No other commercially approved topical formulations of lidocaine (whether creams, lotions or gels) are indicated for neuropathic pain. Non-dermal patch formulations are generally indicated as local anesthetics and anti-pruritics. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. Formulations that do not involve a dermal-patch system are generally indicated as local anesthetics and anti-pruritics. In February 2007 the FDA notified consumers and healthcare professionals of the potential hazards of the use of topical lidocaine. Those at particular risk were individuals that applied large amounts of this substance over large areas, left the products on for long periods of time, or used the agent with occlusive dressings. Systemic exposure was highly variable among patients. Only FDA-approved products are currently recommended. (Argoff, 2006) (Dworkin, 2007) (Khaliq-Cochrane, 2007) (Knotkova, 2007) (Lexi-Comp, 2008) Non-neuropathic pain: Not recommended. There is only one trial that tested 4% lidocaine for treatment of chronic muscle pain. The results showed there was no superiority over placebo. (Scudds, 1995) This medication is recommended for localized peripheral pain. The patient does have lower extremity pain, however the patient has no documented failure of all first line agents indicated for the treatment of neuropathic pain as outlined above. Therefore, criteria as set forth by the California MTUS as outlined above have not been met and the request is not medically necessary.

6 psychologist sessions: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Psychological treatment.

Decision rationale: The California chronic pain medical treatment guidelines section on psychological treatment states: Recommended for appropriately identified patients during treatment for chronic pain. Psychological intervention for chronic pain includes setting goals, determining appropriateness of treatment, conceptualizing a patient's pain beliefs and coping styles, assessing psychological and cognitive function, and addressing co-morbid mood disorders (such as depression, anxiety, panic disorder, and posttraumatic stress disorder). Cognitive behavioral therapy and self-regulatory treatments have been found to be particularly effective. Psychological treatment incorporated into pain treatment has been found to have a positive short-term effect on pain interference and long-term effect on return to work. The following stepped care approach to pain management that involves psychological intervention has been suggested: Step 1: Identify and address specific concerns about pain and enhance interventions that emphasize self-management. The role of the psychologist at this point includes education and training of pain care providers in how to screen for patients that may need early psychological intervention. Step 2: Identify patients who continue to experience pain and disability after the usual time of recovery. At this point a consultation with a psychologist allows for screening,

assessment of goals, and further treatment options, including brief individual or group therapy. Step 3: Pain is sustained in spite of continued therapy (including the above psychological care). Intensive care may be required from mental health professions allowing for a multidisciplinary treatment approach. See also Multi-disciplinary pain programs. See also ODG Cognitive Behavioral Therapy (CBT) Guidelines. (Otis, 2006) (Townsend, 2006) (Kerns, 2005) (Flor, 1992) (Morley, 1999) (Ostelo, 2005) Psychological treatment in particular cognitive behavioral therapy has been found to be particularly effective in the treatment of chronic pain. As this patient has continued ongoing pain, this service is indicated per the California MTUS and thus is medically necessary.