

Case Number:	CM15-0048718		
Date Assigned:	03/20/2015	Date of Injury:	11/06/2012
Decision Date:	05/21/2015	UR Denial Date:	02/17/2015
Priority:	Standard	Application Received:	03/16/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Arizona, Michigan

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 28-year-old female with an industrial injury dated 11/06/2012. Her diagnosis includes neck pain, clinically consistent cervical radiculopathy, myofascial pain and thoracic sprain/strain. Prior management includes diagnostics, physical therapy and medications. She presents on 10/02/2014 with complaints of neck and left upper extremity pain. The injured worker was protective of her left upper extremity. Spasm and tenderness was noted in the cervical paraspinal muscles on the left side. The provider requested medications and pain management consult. The record dated 10/02/2014 is the most recent record submitted. The UR references records dated 2015 however these are not available in the records submitted for this review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 7.5/325 mg #60 with 3 refills: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96 (78,89,95).

Decision rationale: Per the MTUS, opioids should be discontinued if there is no overall improvement in function, unless there are extenuating circumstances, Opioids should be continued if the patient has returned to work or has improved functioning and pain. Ongoing management actions should include prescriptions from a single practitioner, taken as directed and all prescriptions from a single pharmacy. The lowest possible dose should be prescribed to improve pain and function. Documentation should follow the 4 A's of analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors. Long-term users of opioids should be regularly reassessed. In the maintenance phase, the dose should not be lowered if it is working. In addition, patients who receive opioid therapy may sometimes develop unexpected changes in their response to opioids, which includes development of abnormal pain, change in pain pattern, persistence of pain at higher levels than expected. When this happens, opioids can actually increase rather than decrease sensitivity to noxious stimuli. It is important to note that a decrease in opioid efficacy should not always be treated by increasing the dose or adding other opioids, but may actually require weaning. A review of the injured workers medical records that are available to me do not reveal documentation according to the guidelines for ongoing management with opioids and without this information medical necessity is not established. Therefore, this request is not medically necessary.

Mobic 15 mg #60 with 3 refills: Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAID's Page(s): 67-68.

Decision rationale: Per the MTUS, NSAIDs are recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. A review of the injured workers medical records that are available to me reveal subjective and objective documentation of the injured workers pain and the continued use of an NSAID would be appropriate in the injured worker, therefore the request for Mobic15mg # 60 with 3 refills is medically necessary.

Pain management 1 time a month for 4 months to neck and left upper extremity:

Overtured

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Low Back - Lumbar & Thoracic (Acute & Chronic) / office visits.

Decision rationale: Per the MTUS/ ACOEM "Patients whose low back may be work related should receive follow-up care every three to five days by a midlevel practitioner, who can counsel them about avoiding static positions, medication use, activity modification, and other concerns. Take care to answer questions and make these sessions interactive so that patients are fully involved in their recovery. If the patient has returned to work, these interactions may be done on site or by telephone to avoid interfering with modified- or full-work activities. Physician follow-up generally occurs when a release to modified, increased, or full duty is needed, or after appreciable healing or recovery can be expected, on average. Physician follow-up might be expected every four to seven days if the patient is off work and every seven to fourteen days if the patient is working. Per the ODG, office visits are "recommended as determined to be medically necessary. Evaluation and management (E&M) outpatient visits to the offices of medical doctor(s) play a critical role in the proper diagnosis and return to function of an injured worker, and they should be encouraged. The need for a clinical office visit with a health care provider is individualized based upon a review of the patient concerns, signs and symptoms, clinical stability, and reasonable physician judgment. The determination is also based on what medications the patient is taking, since some medicines such as opiates, or medicines such as certain antibiotics, require close monitoring. As patient conditions are extremely varied, a set number of office visits per condition cannot be reasonably established. The determination of necessity for an office visit requires individualized case review and assessment, being ever mindful that the best patient outcomes are achieved with eventual patient independence from the health care system through self care as soon as clinically feasible." Therefore based on the injured workers clinical presentation and the guidelines the request for Pain management 1 time a month for 4 months to neck and left upper extremity is medically necessary.

Butrans 10 mcg #4 with 3 refills: Overtured

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (chronic) / Buprenorphine for chronic pain.

Decision rationale: The MTUS did not specifically address the use of Butrans and therefore other guidelines were consulted. Per the ODG Butrans (buprenorphine) is recommended as an option for treatment of chronic pain in selected patients and is generally not considered first line.

Suggested populations: (1) Patients with a hyperalgesic component to pain; (2) Patients with centrally mediated pain; (3) Patients with neuropathic pain; (4) Patients at high-risk of non-adherence with standard opioid maintenance; (5) For analgesia in patients who have previously been detoxified from other high-dose opioids. Use for pain with formulations other than Butrans is off-label. Due to complexity of induction and treatment, the drug should be reserved for use by clinicians with experience. A review of the injured workers medical records reveal that she falls within the recommended criteria for continued Butrans use, therefore the request for Butrans 10mcg #4 with 3 refills is medically necessary.