

Case Number:	CM15-0063115		
Date Assigned:	04/08/2015	Date of Injury:	10/06/2008
Decision Date:	05/13/2015	UR Denial Date:	03/27/2015
Priority:	Standard	Application Received:	04/02/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 53 year old, male who sustained a work related injury on 10/6/08. He was stepping off an elevator that was not level with floor, his knee locked up, he lost his balance and collapsed. The diagnoses have included osteoarthritis of knee, anterior cruciate ligament tear, medial meniscus tear and status post right knee surgery. Treatments have included a right knee brace, right knee surgery, home exercise program, Flector patches and oral medications. In the PR-2 dated 3/17/15, the injured worker complains of an episode of right knee popping, knee pain and swelling. He describes the pain as dull and occasional. He rates the knee pain a 3-4/10. He states the pain occurs 2-3 times a week. The treatment plan is a refill of Anaprox.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Anaprox DS 550mg #100: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 67-68.

Decision rationale: The MTUS/Chronic Pain Medical Treatment Guidelines comment on the use of NSAIDs, including Anaprox DS, as a treatment modality. The specific recommendations for the use of NSAIDs are as follows: Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP. For patients with acute low back pain with sciatica a recent Cochrane review (including three heterogeneous randomized controlled trials) found no differences in treatment with NSAIDs vs. placebo. In patients with axial low back pain this same review found that NSAIDs were not more effective than acetaminophen for acute low-back pain, and that acetaminophen had fewer side effects. Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. In this case, the records indicate that Anaprox DS is being used as a long-term treatment strategy for this patient's chronic pain. Per the above cited MTUS guidelines, NSAIDs such as Anaprox DS are recommended for short-term symptomatic relief. The use of Anaprox DS extends well beyond these MTUS recommendations. For this reason, Anaprox DS is not considered as medically necessary.