

Case Number:	CM15-0063841		
Date Assigned:	04/09/2015	Date of Injury:	09/17/1991
Decision Date:	05/12/2015	UR Denial Date:	03/05/2015
Priority:	Standard	Application Received:	04/03/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials: State(s) of Licensure: New Jersey, Michigan, California Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 72-year-old male, who sustained an industrial injury on September 17, 1991. He reported back pain and bilateral knee pain. The injured worker was diagnosed as having lumbar facet arthrosis, lumbar spinal stenosis, thoracic pain with radicular symptoms around the ribs, low back pain and bilateral knee pain. Treatment to date has included radiographic imaging, diagnostic studies, a pain generator, conservative care, medications and activity modifications. Currently, the injured worker complains of mid and low back pain radiating to the thighs and bilateral knee pain. The injured worker reported an industrial injury in 1991, resulting in the above noted pain. He was treated conservatively without complete resolution of the pain. Evaluation on April 10, 2015, revealed continued chronic pain. He reported needing a refill of medications. He has been on a chronic pain management regiment with pain medications for quite some time to remain functional. Medications were requested.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg quantity 180: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, Norco (Hydrocodone/Acetaminophen) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. According to the patient file, there is no objective documentation of pain and functional improvement to justify continuous use of Norco. Norco was used for longtime without documentation of functional improvement or evidence of improvement of activity of daily living. Therefore, the prescription of Norco 10/325 mg # 180 is not medically necessary.

Restoril 15mg quantity 60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

Decision rationale: According to ODG guidelines and in the treatment of insomnia section. Recommend that treatment be based on the etiology, with the medications recommended below. See Insomnia. Pharmacological agents should only be used after careful evaluation of potential causes of sleep disturbance. Failure of sleep disturbance to resolve in a 7 to 10 day period may indicate a psychiatric and/or medical illness. (Lexi-Comp, 2008) Primary insomnia is generally addressed pharmacologically. Secondary insomnia may be treated with pharmacological and/or psychological measures. The specific component of insomnia should be addressed: (a) Sleep onset; (b) Sleep maintenance; (c) Sleep quality; & (d) Next-day functioning. Pharmacologic Treatment: There are four main categories of pharmacologic treatment: (1) Benzodiazepines; (2) Non-benzodiazepines; (3) Melatonin & melatonin receptor agonists; & (4) Over-the-counter medications. The majority of studies have only evaluated short-term treatment (i.e., 4 weeks) of insomnia; therefore, more studies are necessary to evaluate the efficacy and safety of treatments for long-term treatment of insomnia. In 2007, the FDA requested that manufacturers of all sedative-hypnotic drugs strengthen product labeling regarding risks (i.e., severe allergic reactions and complex sleep-related behaviors, such as sleep driving). It is recommended that treatments for insomnia should reduce time to sleep onset, improve sleep maintenance, avoid residual effects and increase next-day functioning. (Morin, 2007) (Reeder, 2007) (1) Benzodiazepines: FDA-approved benzodiazepines for sleep maintenance insomnia include estazolam (ProSom), flurazepam (Dalmane), quazepam (Doral), and temazepam (Restoril). Triazolam (Halcion) is FDA-approved for sleep-onset insomnia. These medications are only recommended for short-term use due to risk of tolerance, dependence, and adverse events (daytime drowsiness, anterograde amnesia, next-day sedation, impaired cognition, impaired

psychomotor function, and rebound insomnia). These drugs have been associated with sleep-related activities such as sleep driving, cooking and eating food, and making phone calls (all while asleep). Particular concern is noted for patients at risk for abuse or addiction.

Withdrawal occurs with abrupt discontinuation or large decreases in dose. Decrease slowly and monitor for withdrawal symptoms. Benzodiazepines are similar in efficacy to benzodiazepine-receptor agonists; however, the less desirable side-effect profile limits their use as a first-line agent, particularly for long-term use. According to MTUS guidelines, benzodiazepines are not recommended for long term use for pain management because of unproven long term efficacy and because of the risk of dependence. Most guidelines limit their use to 4 weeks. There is no recent documentation of insomnia related to pain. Therefore the prescription of Restoril 15mg quantity 60 is not medically necessary.

Lyrica 75mg quantity 60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Lyrica; Anti Epilepsy Drugs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Lyrica Page(s): 20.

Decision rationale: According to MTUS guidelines, Lyrica is an anti-epilepsy drug (AEDs - also referred to as anti-convulsant), which has been shown to be effective for treatment of diabetic; painful neuropathy and post-therapeutic neuralgia; and has been considered as a first-line treatment for neuropathic pain. There is no clear documentation of neuropathic pain in this patient that required and responded to previous use of Lyrica. In addition, there is no clear proven efficacy of Lyrica for back pain. Therefore, Lyrica 50mg #60 is not medically necessary.