

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM15-0064130 | | |
| Date Assigned: | 04/09/2015 | Date of Injury: | 10/05/2012 |
| Decision Date: | 05/08/2015 | UR Denial Date: | 03/25/2015 |
| Priority: | Standard | Application Received: | 04/03/2015 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Physical Medicine & Rehabilitation

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 39 year old male, who sustained an industrial injury on October 5, 2012. He has reported neck pain, back pain, wrist pain, and elbow pain. Diagnoses have included degenerative lumbar spondylosis, myofascial pain syndrome, chronic right shoulder pain secondary to osteoarthritis, right epicondylitis, carpal tunnel syndrome, insomnia, and degenerative cervical spondylosis. Treatment to date has included medications, physical therapy, right elbow surgery, and imaging studies. A progress note dated February 17, 2015 indicates a chief complaint of neck pain, back pain, right side pain, and increased leg pain. The treating physician documented a plan of care that included medications.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Oxycodone/APAP Tablet 10/325 mg, Qty 120: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

Decision rationale: Per the MTUS Guidelines cited, opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury without acute flare, new injury, or progressive deterioration. The Oxycodone/APAP Tablet 10/325 mg, Qty 120 is not medically necessary and appropriate.

Diazepam Tablet 10 mg, Qty 30: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines, page 23.

Decision rationale: Valium (Diazepam) is an anti-anxiety medication in the benzodiazepine family and like other benzodiazepines, act by enhancing the effects of gamma-aminobutyric acid (GABA) in the brain. GABA is a neurotransmitter (a chemical that nerve cells use to communicate with each other) which inhibits many of the activities of the brain. It is believed that excessive activity in the brain may lead to anxiety or other psychiatric disorders. Valium also is used to prevent certain types of seizures. Valium is used for the short-term relief of the symptoms of anxiety. It is used for certain types of seizures, specifically petit mal seizures, akinetic seizures, and myoclonus, as well as Lennox-Gastaut syndrome. Submitted reports have not adequately addressed the indication for Valium's continued use for the chronic injury. Per the Chronic Pain Treatment Guidelines, benzodiazepines are not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks as chronic benzodiazepines are the treatment of choice in very few conditions and tolerance to hypnotic effects develops rapidly. Additionally, submitted reports have not demonstrated clear functional benefit of treatment already rendered. The Diazepam Tablet 10 mg, Qty 30 is not medically necessary and appropriate.

Lunesta Tablet 3 mg, Qty 60: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Insomnia Treatment, pages 535-536.

Decision rationale: Hypnotics are not included among the multiple medications noted to be optional adjuvant medications, per the Official Disability Guidelines (ODG), Pain. Additionally, Lunesta is a non-benzodiazepine-like, Schedule IV controlled substance. Long-term use is not recommended as efficacy is unproven with a risk of dependence. Most guidelines limit use to 4 weeks. Their range of action includes sedative/hypnotic and anxiolytic. Chronic use is the treatment of choice in very few conditions. Tolerance to hypnotic effects develops rapidly. Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. Submitted documents have not demonstrated any functional improvement from treatment rendered for this chronic injury. The Lunesta Tablet 3 mg, Qty 60 is not medically necessary and appropriate.