

Case Number:	CM15-0064899		
Date Assigned:	04/13/2015	Date of Injury:	09/11/2014
Decision Date:	05/15/2015	UR Denial Date:	04/01/2015
Priority:	Standard	Application Received:	04/06/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: New York, Tennessee
Certification(s)/Specialty: Emergency Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 34-year-old male, who sustained an industrial injury on 9/11/14. The injured worker has complaints of low back pain and right leg pain. The diagnoses have included lumbar discogenic disease L5-S1. Treatment to date has included physical therapy; trigger point injection; lumbar spine X-rays; magnetic resonance imaging (MRI); activity modification, prolonged rest and medications. The documentation noted that the radicular pain is due to lumbar spinal stenosis. The request was for bilateral L4-S1 epidural steroid injection times three.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Bilateral L4-S1 ESI x3: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 46.

Decision rationale: Epidural steroid injections are recommended as an option for treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of

radiculopathy). Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. Epidural steroid injection can offer short term pain relief and use should be in conjunction with other rehab efforts, including continuing a home exercise program. There is little information on improved function. The American Academy of Neurology recently concluded that epidural steroid injections may lead to an improvement in radicular lumbosacral pain between 2 and 6 weeks following the injection, but they do not affect impairment of function or the need for surgery and do not provide long-term pain relief beyond 3 months, and there is insufficient evidence to make any recommendation for the use of epidural steroid injections to treat radicular cervical pain. Criteria for the use of Epidural steroid injections: Note: The purpose of ESI is to reduce pain and inflammation, thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit. 1) Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. 2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants). 3) Injections should be performed using fluoroscopy (live x-ray) for guidance. 4) If used for diagnostic purposes, a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. Diagnostic blocks should be at an interval of at least one to two weeks between injections. 5) No more than two nerve root levels should be injected using transforaminal blocks. 6) No more than one interlaminar level should be injected at one session. 7) In the therapeutic phase, repeat blocks should be based on continued objective documented pain and functional improvement, including at least 50% pain relief with associated reduction of medication use for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year. 8) Current research does not support a "series-of-three" injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections. 9) Epidural steroid injection is not to be performed on the same day as trigger point injection, sacroiliac joint injection, facet joint injection or medial branch block. In this case, the request was for a series of three injections. Per criteria, this is not supported by current research. The request is not medically necessary.