

Case Number:	CM15-0065357		
Date Assigned:	04/13/2015	Date of Injury:	10/19/1999
Decision Date:	05/12/2015	UR Denial Date:	03/26/2015
Priority:	Standard	Application Received:	04/07/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 60 year old female, who sustained an industrial injury on October 19, 1999. The injured worker was diagnosed as having chronic low back pain, lumbar radiculopathy, status post four lumbar spine surgeries, chronic pain syndrome, and failed back surgery syndrome. Treatment to date has included multiple lumbar epidural injections, lumbar spine MRI, lumbar surgeries, and medication. Currently, the injured worker complains of low back and right lower extremity pain. The Primary Treating Physician's report dated February 12, 2015, noted the injured worker reporting that she was doing worse, with the low back pain rated n 8/10 on the pain scale. The injured worker was noted to be taking Oxycontin, Vicoprofen, and Gabapentin. Physical examination was noted to show the lumbar spine with decreased flexion and extension, decreased sensation in the right L4, L5, and S1 dermatomes, and muscle stretch reflex diminished but sympathetic at the patella and Achilles bilaterally. The lumbar spine MRI dated December 11, 2012, was noted to show facet arthropathy, L4-L5 severe left neural foraminal narrowing and moderate right neural foraminal narrowing. The treatment plan was noted to include requests for authorization for medications including prescriptions for Gabapentin, MS Contin, and Norco, follow up pain management in four weeks, and pain psych evaluation and clearance as prerequisite for spinal cord stimulator trial.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MS Contin 30mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug- related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. There is no clear documentation of patient improvement in level of function, quality of life, adequate follow up for absence of side effects and aberrant behavior with a previous use of narcotics. The patient continues to have chronic pain despite the continuous use of narcotics. Therefore, the request for MS Contin 30mg #60 is not medically necessary.

Norco 7.5/325mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, Norco (Hydrocodone/Acetaminophen) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug- related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. According to the patient file, there is no objective documentation of pain and functional improvement to justify continuous use of Norco. Norco was used for longtime without documentation of functional improvement or evidence of return to work or improvement of activity of daily living. Therefore, the prescription of Norco 7.5/325mg #90 is not medically necessary.