

<b>Case Number:</b>	CM15-0065597		
<b>Date Assigned:</b>	04/13/2015	<b>Date of Injury:</b>	01/12/1998
<b>Decision Date:</b>	06/11/2015	<b>UR Denial Date:</b>	03/19/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	04/07/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials: State(s) of Licensure: North Carolina Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 63 year old female who sustained an industrial injury on January 12, 1998. Prior treatment includes Synvisc injection, right knee lateral meniscectomy, chondroplasty and medications. Currently the injured worker complains of pain in the knee. Diagnoses associated with the request include pain in the joint of the lower leg. The treatment plan includes medications of Flexeril, Lidoderm and Lunesta.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Flexeril 10 mg, sixty count:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 64.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines muscle relaxants Page(s): 63-65.

**Decision rationale:** The California chronic pain medical treatment guidelines section on muscle relaxants states: Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder,

2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) (Chou, 2004) This medication is not intended for long-term use per the California MTUS. The medication has not been prescribed for the flare-up of chronic low back pain. This is not an approved use for the medication. For these reasons, criteria for the use of this medication have not been met. Therefore the request is not medically necessary.

**Lidoderm patch 5%, ninety count:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 111 - 112.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines topical analgesics Page(s): 111-113.

**Decision rationale:** The California chronic pain medical treatment guidelines section on topical lidocaine states: Lidocaine Indication: Neuropathic pain Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Topical lidocaine, in the formulation of a dermal patch (Lidoderm) has been designated for orphan status by the FDA for neuropathic pain. Lidoderm is also used off-label for diabetic neuropathy. No other commercially approved topical formulations of lidocaine (whether creams, lotions or gels) are indicated for neuropathic pain. Non-dermal patch formulations are generally indicated as local anesthetics and anti-pruritics. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. Formulations that do not involve a dermal-patch system are generally indicated as local anesthetics and anti-pruritics. In February 2007 the FDA notified consumers and healthcare professionals of the potential hazards of the use of topical lidocaine. Those at particular risk were individuals that applied large amounts of this substance over large areas, left the products on for long periods of time, or used the agent with occlusive dressings. Systemic exposure was highly variable among patients. Only FDA-approved products are currently recommended. (Argoff, 2006) (Dworkin, 2007) (Khaliq-Cochrane, 2007) (Knotkova, 2007) (Lexi-Comp, 2008) Non-neuropathic pain: Not recommended. There is only one trial that tested 4% lidocaine for treatment of chronic muscle pain. The results showed there was no superiority over placebo. (Scudds, 1995) This medication is recommended for localized peripheral pain. The patient has no documented failure of all first line agents indicated for the treatment of neuropathic pain as outlined above. Therefore criteria as set forth by the California MTUS as outlined above have not been met and the request is not medically necessary.

**Lunesta 2 mg, sixty count:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter.

**MAXIMUS guideline:** The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation ODG, insomnia.

**Decision rationale:** The California MTUS and the ACOEM do not specifically address this medication. Per the official disability guidelines recommend pharmacological agents for insomnia only is used after careful evaluation of potential causes of sleep disturbance. Primary insomnia is usually addressed pharmacologically. Secondary insomnia may be treated with pharmacological and/or psychological measures. Pharmacological treatment consists of four main categories: Benzodiazepines, Non-benzodiazepines, Melatonin and melatonin receptor agonists and over the counter medications. Sedating antidepressants have also been used to treat insomnia however there is less evidence to support their use for insomnia, but they may be an option in patients with coexisting depression. The patient does not have the diagnosis of primary insomnia nor is there provided clinical documentation demonstrating failure of sleep hygiene measures/counseling. Therefore the request is not medically necessary.