

Case Number:	CM15-0066528		
Date Assigned:	04/14/2015	Date of Injury:	11/19/2012
Decision Date:	05/13/2015	UR Denial Date:	03/17/2015
Priority:	Standard	Application Received:	04/07/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Arizona, California

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 59-year-old female, who sustained an industrial injury on 11/19/12. She reported low back, left hip, left ankle and left knee injury. The injured worker was diagnosed as having facet joint hypertrophy at L3-4, L4-5 and L5-S1, left lower extremity radiculopathy, osteoarthritis of left hip and anterior talofibular ligament tear of left ankle. Treatment to date has included oral medications including opioids, physical therapy, acupuncture, and medial branch blocks. Currently, the injured worker complains of left knee, left ankle, left hip and left sided low back pain. The injured worker states the medications helps keep her pain at a tolerable level. Physical exam noted tenderness and guarding in lumbar paraspinal musculature with decreased range of motion. The treatment plan included an epidural injection at L5-S1.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Retrospective request for anes nerve blocks & injection (DOS: 1/5/15): Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Low Back, Faet joint diagnostic blocks (injections); Pain, Epidural steroid injections (ESIs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines epidural injections Page(s): 47. Decision based on Non-MTUS Citation ODG- Back pain and pg 46.

Decision rationale: According to the guidelines, Criteria for the use of diagnostic blocks for facet 'mediated' pain: Clinical presentation should be consistent with facet joint pain, signs & symptoms. 1. One set of diagnostic medial branch blocks is required with a response of 70%. The pain response should last at least 2 hours for Lidocaine. 2. Limited to patients with low-back pain that is non-radicular and at no more than two levels bilaterally. 3. There is documentation of failure of conservative treatment (including home exercise, PT and NSAIDs) prior to the procedure for at least 4-6 weeks. 4. No more than 2 facet joint levels are injected in one session (see above for medial branch block levels). 5. Recommended volume of no more than 0.5 cc of injectate is given to each joint. 6. No pain medication from home should be taken for at least 4 hours prior to the diagnostic block and/or 4 to 6 hours afterward. 7. Opioids should not be given as a 'sedative' during the procedure. 8. The use of IV sedation (including other agents such as midazolam) may be grounds to negate the results of a diagnostic block, and should only be given in cases of extreme anxiety. 9. The patient should document pain relief with an instrument such as a VAS scale, emphasizing the importance of recording the maximum pain relief and maximum duration of pain. The patient should also keep medication use and activity logs to support subjective reports of better pain control. 10. Diagnostic facet blocks should not be performed in patients in whom a surgical procedure is anticipated. (Resnick, 2005) 11. Diagnostic facet blocks should not be performed in patients who have had a previous fusion procedure at the planned injection level. According to the guidelines, the criteria for the use of Epidural steroid injections: Note: The purpose of ESI is to reduce pain and inflammation, restoring range of motion and thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit. 1) Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. 2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants). 3) Injections should be performed using fluoroscopy (live x-ray) for guidance. 4) If used for diagnostic purposes, a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. Diagnostic blocks should be at an interval of at least one to two weeks between injections. 5) No more than two nerve root levels should be injected using transforaminal blocks. 6) No more than one interlaminar level should be injected at one session. 7) In the therapeutic phase, repeat blocks should be based on continued objective documented pain and functional improvement, including at least 50% pain relief with associated reduction of medication use for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year. (Manchikanti, 2003) (CMS, 2004) (Boswell, 2007) 8) Current researches do not support a 'series-of-three' injection in either the diagnostic or the therapeutic phase. We recommend no more than 2 ESI injections. In this case, the claimant has had an MRI with no significant foraminal narrowing or radicular component on the left side. EMG showed no radiculopathy in the lumbar spine but bilaterally in the S1 level. The area of injection was not specified. There was no plan for a facer neurotomy. The request for the block and injection above is not medically necessary.