

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM15-0066873 | | |
| Date Assigned: | 04/14/2015 | Date of Injury: | 09/15/2008 |
| Decision Date: | 06/03/2015 | UR Denial Date: | 04/02/2015 |
| Priority: | Standard | Application Received: | 04/08/2015 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: New York

Certification(s)/Specialty: Anesthesiology

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 31-year-old female, who sustained an industrial injury on 9/15/2008. She reported injury when a patient fell on her. The injured worker was diagnosed as having cervical sprain and lumbar sprain. Recent diagnostic study was not documented in the medical records. Treatment to date has included medication management. In a progress note dated 3/25/2015, the injured worker complains of right shoulder pain and insomnia. The treating physician is requesting Tramadol, Cyclobenzaprine, Terocin patches, Exoten-C lotion, Fenoprofen and Omeprazole.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Tramadol extended release 150mg quantity 30 with three refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids for the treatment of chronic pain Page(s): 91-97.

Decision rationale: According to the California MTUS, Tramadol (Ultram) is a synthetic opioid, which affects the central nervous system and is indicated for the treatment of moderate to severe pain. Per CA MTUS Guidelines, certain criteria need to be followed, including an ongoing review and documentation of pain relief and functional status, appropriate medication use, and side effects. Pain assessment should include current pain: last reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid, and the duration of pain relief. According to the medical records, there has been no objective evidence of this medication's analgesic effectiveness or functional improvement, and no clear documentation that the patient has responded to ongoing opioid therapy. Medical necessity of the requested medication has not been established. Of note, discontinuation of an opioid analgesic requires a taper to avoid withdrawal symptoms. The requested Tramadol ER with (3) refills is not medically necessary.

Cyclobenzaprine 7.5 mg quantity 60 with three refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 63.

Decision rationale: According to the reviewed literature, Cyclobenzaprine (Flexeril) is not recommended for the long-term treatment of chronic pain. The medication has its greatest effect in the first four days of treatment. Guidelines state that this medication is not recommended to be used for longer than 2-3 weeks. According to CA MTUS Guidelines, muscle relaxants are not considered any more effective than non-steroidal anti-inflammatory medications alone. There is no documentation of functional improvement from any previous use of this medication. Based on the currently available information, the medical necessity for this muscle relaxant medication has not been established. The requested medication, with (3) refills, is not medically necessary.

Terocin Patches quantity 30 with three refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

Decision rationale: According to the California MTUS Guidelines, topical analgesics are primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. These agents are applied topically to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. Many agents are compounded as monotherapy or in combination for pain control including, for example, NSAIDs, opioids, capsaicin, local anesthetics or antidepressants. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. In this case, there is no documentation provided necessitating Terocin. This medication contains methyl

salicylate, capsaicin, menthol, and lidocaine. MTUS states that capsaicin is recommended only as an option in patients who have not responded or are intolerant to other treatments. There is no documentation of intolerance to other previous oral medications. Medical necessity for the requested topical medication has not been established. The requested Terocin patches with (3) refills are not medically necessary.

Exoten-C Lotion 120gm quantity 2 with three refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

Decision rationale: According to the California MTUS Guidelines, topical analgesics are primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. These agents are applied topically to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. Many agents are compounded as monotherapy or in combination for pain control including, for example, NSAIDs, opioids, capsaicin, local anesthetics or antidepressants. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. In this case, Exoten-C Lotion contains methyl salicylate, menthol, and capsaicin. MTUS states that capsaicin is recommended only as an option in patients who have not responded or are intolerant to other treatments. There is no documentation of intolerance to other previous oral medications. Medical necessity for the requested topical medication has not been established. The requested treatment, with (3) refills, is not medically necessary.

Fenoprofen 400mg quantity 60 with three refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 21, 67-71. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) NSAIDs.

Decision rationale: Fenoprofen calcium (Nalfon) is a non-steroidal anti-inflammatory drug (NSAID). Oral NSAIDs are recommended for the treatment of chronic pain and control of inflammation as a second-line therapy after acetaminophen. According to the California MTUS Guidelines, NSAIDs reduce pain so activity and functional restoration can resume, but long-term use may not be warranted. ODG states that NSAIDs are recommended for acute pain, acute low back pain (LBP), short-term pain relief in chronic LBP, and short-term improvement of function in chronic LBP. There is no evidence of long-term effectiveness for pain or function. Current evidence based guidelines indicate that Fenoprofen is an NSAID medication which is less effective, and has greater side effects than Naproxen or Ibuprofen. Guidelines indicate that Fenoprofen should not be used unless there is a sound medical basis for not using a safer or more

effective alternative NSAID. In this case, there was no rationale provided which explained the request for Fenoprofen. Medical necessity of the requested medication has not been established. The requested medication with (3) refills is not medically necessary.

Omeprazole 20mg quantity 60 with three refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines PPIs Page(s): 68. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) PPIs.

Decision rationale: According to CA MTUS (2009), proton pump inhibitors, such as Omeprazole (Prilosec), are recommended for patients taking NSAIDs with documented GI distress symptoms or specific GI risk factors. There is no documentation indicating the patient has any GI symptoms or GI risk factors. Risk factors include, age >65, history of peptic ulcer disease, GI bleeding, concurrent use of aspirin, corticosteroids, and/or anticoagulants or high-dose/multiple NSAIDs. There is no documentation of any reported GI complaints. Based on the available information provided for review, the medical necessity for Prilosec has not been established. The requested medication, with (3) refills, is not medically necessary.