

Case Number:	CM15-0068619		
Date Assigned:	04/16/2015	Date of Injury:	09/04/2009
Decision Date:	05/15/2015	UR Denial Date:	03/18/2015
Priority:	Standard	Application Received:	04/10/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Indiana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 62 year old male, who sustained an industrial injury on September 4, 2009, incurring neck and shoulder injuries. He was diagnosed with cervical degenerative disc disease and cervical stenosis and chronic shoulder pain. Treatment included home exercise program, acupuncture, epidural steroid injection anti-inflammatory drugs and pain medications. Currently, the injured worker complained of persistent neck pain and left upper extremity pain. The treatment plan that was requested for authorization included prescriptions for Norco and Zanaflex.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids for chronic pain Page(s): 80.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG)

Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for neck and low back pain "except for short use for severe cases, not to exceed 2 weeks." The requested treatment is for more than 2 weeks. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for Norco is not medically necessary.

Zanaflex 4mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 64-66.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants; Zanaflex Page(s): 63-66.

Decision rationale: Zanaflex is the brand name version of tizanidine, which is a muscle relaxant. MTUS states concerning muscle relaxants "Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (VanTulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also, there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) Sedation is the most commonly reported adverse effect of muscle relaxant medications. These drugs should be used with caution in patients driving motor vehicles or operating heavy machinery. Drugs with the most limited published evidence in terms of clinical effectiveness include chlorzoxazone, methocarbamol, dantrolene and baclofen. (Chou, 2004) According to a recent review in American Family Physician, skeletal muscle relaxants are the most widely prescribed drug class for musculoskeletal conditions (18.5% of prescriptions), and the most commonly prescribed antispasmodic agents are carisoprodol, cyclobenzaprine, metaxalone, and methocarbamol, but despite their popularity, skeletal muscle relaxants should not be the primary drug class of choice for musculoskeletal conditions (See2, 2008)." MTUS further states, "Tizanidine (Zanaflex, generic available) is a centrally acting alpha2-adrenergic agonist that is FDA approved for management of spasticity; unlabeled use for low back pain. (Malanga, 2008) Eight studies have demonstrated efficacy for low back pain. (Chou, 2007) One study (conducted only in females) demonstrated a significant decrease in pain associated with chronic myofascial pain syndrome and the authors recommended its use as a first line option to treat myofascial pain. (Malanga,

2002) May also provide benefit as an adjunct treatment for fibromyalgia (ICSI, 2007)." There is no documentation of failure of first line therapies. The requested quantity is beyond the short duration that the guidelines recommend. There is no documentation of an acute exacerbation of low back pain. Therefore, the request is not medically necessary.