

Case Number:	CM15-0069041		
Date Assigned:	04/16/2015	Date of Injury:	05/22/2009
Decision Date:	05/19/2015	UR Denial Date:	04/13/2015
Priority:	Standard	Application Received:	04/13/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 47 year old female who sustained an industrial injury on 5/22/09. The diagnoses have included lumbar post laminectomy syndrome, low back pain and chronic pain syndrome. Treatment to date has included medications, diagnostics, activity modifications and physical therapy. Currently, as per the physician progress note dated 3/5/15, the injured worker complains of pain in the back with significant coping issues. The physical exam was unchanged. There was tenderness in the lumbar spine and painful range of motion. The treatment plans were for follow up in 4 weeks and renew medications. The physician requested treatments included Tramadol 50mg #90, Butrans 10mcg #4 and Flexeril 7.5mg #90.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Tramadol 50mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Page(s): 93-94.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Page(s): 113.

Decision rationale: According to MTUS guidelines, Ultram (Tramadol) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. There is no clear documentation of pain and functional improvement with previous use of Tramadol. There is no clear documentation of continuous monitoring of patient's compliance with her medications. There is no documentation of the medical necessity of Tramadol over NSAID. Therefore, the prescription of Tramadol 50mg #90 is not medically necessary.

Butrans 10mcg #4: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Buprenorphine Page(s): 26-27.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 78-79.

Decision rationale: According to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side

effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. According to MTUS guidelines, Butrans is recommended to treat opiate addiction. There is no clear documentation of patient improvement in level of function, quality of life, adequate follow up or absence of side effects and aberrant behavior with previous use of opioids. The patient continued to have significant pain with Butrans. There is no justification to use multiple opioids. There is no recent documentation of recent opioid addiction. Therefore, the request for Butrans 10mcg #4 is not medically necessary.

Flexeril 7.5mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine (Flexeril) Page(s): 41.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 63.

Decision rationale: According to MTUS guidelines, Flexeril, a non-sedating muscle relaxants, is recommended with caution as a second line option for short-term treatment of acute exacerbations in patients with chronic spasm and pain. Efficacy appears to diminish over time and prolonged use may cause dependence. There is no recent evidence of pain flare or spasm relief with the previous use of Flexeril. Therefore the request for Flexeril 7.5mg #90 is not medically necessary.