

Case Number:	CM15-0070366		
Date Assigned:	04/20/2015	Date of Injury:	03/05/2001
Decision Date:	05/22/2015	UR Denial Date:	03/18/2015
Priority:	Standard	Application Received:	04/14/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Internal Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 58-year-old male who sustained an industrial injury on March 5, 2001. He has reported lower back pain and left leg pain and has been diagnosed with chronic pain syndrome and lumbar post laminectomy syndrome, lumbar region. Treatment has included medication, lying down, ice, and external stimulation. Currently the injured worker complains of severe back pain 8/10 with weakness and numbness to the left leg. The treatment plan included Norco and Avinza.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg, #120: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, criteria for use Page(s): 76-78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page 74-96. Hydrocodone/Acetaminophen Page 91.

Decision rationale: Medical Treatment Utilization Schedule (MTUS) Chronic Pain Medical Treatment Guidelines (page 89) present the strategy for maintenance for long-term users of opioids. "Do not attempt to lower the dose if it is working." Supplemental doses of break-through medication may be required for incidental pain, end-of dose pain, and pain that occurs with predictable situations. The standard increase in dose is 25 to 50% for mild pain and 50 to 100% for severe pain. Actual maximum safe dose will be patient-specific and dependent on current and previous opioid exposure, as well as on whether the patient is using such medications chronically. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant drug-related behaviors. These domains have been summarized as the 4 A's (analgesia, activities of daily living, adverse side effects, and aberrant drug-taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. Hydrocodone/Acetaminophen (Norco) is indicated for moderate to moderately severe pain. The treating physician's progress report dated 3/05/15 documented a history of lumbar fusion spine surgery, knee surgery, spinal cord stimulator implantation, and lumbar post laminectomy syndrome. Analgesia, activities of daily living, adverse side effects, and aberrant behaviors were addressed. Medical records document objective physical examination findings. Medical records document regular physician clinical evaluations and monitoring. Per MTUS, Hydrocodone/Acetaminophen (Norco) is indicated for moderate to moderately severe pain. The request for Norco (Hydrocodone/Acetaminophen) is supported by the MTUS guidelines. Therefore, the request for Norco 10/325 mg is medically necessary.

Avinza 120mg, #60: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, criteria for use Page(s): 76-78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page 74-96.

Decision rationale: Medical Treatment Utilization Schedule (MTUS) Chronic Pain Medical Treatment Guidelines (page 89) present the strategy for maintenance for long-term users of opioids. Do not attempt to lower the dose if it is working. Supplemental doses of break-through medication may be required for incidental pain, end-of dose pain, and pain that occurs with predictable situations. The standard increase in dose is 25 to 50% for mild pain and 50 to 100% for severe pain. Actual maximum safe dose will be patient-specific and dependent on current and previous opioid exposure, as well as on whether the patient is using such medications chronically. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant drug-related behaviors. These domains have been summarized as the 4 A's (analgesia, activities of daily living, adverse side effects, and aberrant drug-taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. The treating physician's progress report dated 3/05/15 documented a history of lumbar fusion spine surgery, knee surgery, spinal cord stimulator implantation, and lumbar post

laminectomy syndrome. Analgesia, activities of daily living, adverse side effects, and aberrant behaviors were addressed. Medical records document objective physical examination findings. Medical records document regular physician clinical evaluations and monitoring. The request for Avinza is supported by the MTUS guidelines. Therefore, the request for Avinza is medically necessary.