

Case Number:	CM15-0071345		
Date Assigned:	04/21/2015	Date of Injury:	10/17/2005
Decision Date:	05/19/2015	UR Denial Date:	03/13/2015
Priority:	Standard	Application Received:	04/14/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Physical Medicine & Rehabilitation

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is an 80 year old male, who sustained an industrial injury on October 17, 2005. He reported burning back pain radiating into the left lower leg. The injured worker was diagnosed as having lumbar intervertebral disc displacement and thoracic or lumbosacral neuritis or radiculitis unspecified. Treatment to date has included work modifications and opioid medication. On March 5, 2015, the injured worker complains of slight worsening of his back pain with radiation down the posterior left leg to the foot. His pain is described as moderate to severe aching. On average he takes his opioid medication 3 per day for pain not relieved by other medications. The physical exam revealed inability to stand erect, an antalgic gait, reversal of the lumbar lordosis, moderate pain to palpation of the bilateral paraspinal muscles of the low back, slight to moderate muscle spasm, slightly restricted range of motion due to pain. The left leg reflexes were absent. There was normal muscle strength and sensation of the lower extremities. The treatment plan includes opioid medication.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325 mg, ninety count: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

Decision rationale: Per the MTUS Guidelines cited, opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury without acute flare, new injury, or progressive deterioration. The Norco 10/325 mg, ninety count is not medically necessary and appropriate.