

Case Number:	CM15-0074279		
Date Assigned:	04/24/2015	Date of Injury:	07/14/2009
Decision Date:	05/21/2015	UR Denial Date:	03/30/2015
Priority:	Standard	Application Received:	04/20/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 49 year old female who sustained an industrial injury on 7/14/09. The injured worker reported symptoms in the bilateral upper extremities. The injured worker was diagnosed as having right lateral epicondylitis, status post left carpal tunnel syndrome, left de Quervain's disease, status post right carpal tunnel, right de Quervain's disease, status post left de Quervain's release. Treatments to date have included status post bilateral hand surgery, Electromyography, Nerve Conduction Velocity studies, and non-steroidal anti-inflammatory drugs. Currently, the injured worker complains of bilateral hand weakness. The plan of care was for medication prescriptions and a follow up appointment at a later date.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Ibuprofen 800mg: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Ibuprofen, NSAIDs Page(s): 67-72.

Decision rationale: MTUS recommends the use of NSAIDs for the acute exacerbation of pain at the lowest effective dose for the shortest amount of time due to the increased cardiovascular risk, renal, hepatic and GI side effects associated with long term use. MTUS states Ibuprofen (Motrin, Advil [otc], generic available): 300, 400, 600, 800 mg. Dosing: Osteoarthritis and off-label for ankylosing spondylitis: 1200 mg to 3200 mg daily. Individual patients may show no better response to 3200 mg as 2400 mg, and sufficient clinical improvement should be observed to offset potential risk of treatment with the increased dose. Higher doses are generally recommended for rheumatoid arthritis: 400-800 mg PO 3-4 times a day, use the lowest effective dose. Higher doses are usually necessary for osteoarthritis. Doses should not exceed 3200 mg/day. Mild pain to moderate pain: 400 mg PO every 4-6 hours as needed. Doses greater than 400 mg have not provided greater relief of pain?. The patient reported on going GI symptoms with concurrent use of Omeprazole while taking Ibuprofen. The treating physician did not document a decrease in pain or functional improvement from the use of Ibuprofen. As such the request for Ibuprofen 800mg is not medically necessary.

Omeprazole 20mg: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI Symptoms and Cardiovascular Risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: MTUS and ODG states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease:(1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or(2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." The medical documents provided do not establish the patient has having documented GI bleeding, perforation, peptic ulcer, high dose NSAID, or other GI risk factors as outlined in MTUS. As such, the request for Omeprazole 20mg is not medically necessary.