

Case Number:	CM15-0074316		
Date Assigned:	04/24/2015	Date of Injury:	12/28/2003
Decision Date:	05/22/2015	UR Denial Date:	04/14/2015
Priority:	Standard	Application Received:	04/20/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 46-year-old male, who sustained an industrial injury on 12/28/03. He reported low back pain with radiation to bilateral legs. The injured worker was diagnosed as having lumbar facet syndrome, lumbar radiculopathy, low back pain, and spondylolisthesis. Treatment to date has included a functional restoration program and medications. A physician's report dated 8/25/14 noted the injured worker was taking Flexeril 10mg and Gralise 600mg. A physician's report dated 3/25/15 noted physical examination findings of paravertebral muscles spasms, tenderness, and tight muscle bands bilaterally on palpation. The injured worker rated his pain as 6/10 with medication and 10/10 without medication. Currently, the injured worker complains of lower back pain with bilateral lower extremity pain. The treating physician requested authorization for Gralise ER 600mg #120 and Flexeril 10mg #30.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Gralise ER: 600mg #120: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 18-19.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-Epilepsy Drugs Page(s): 16-22. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, Anti-epilepsy drugs (AEDs) for pain, Gabapentin (Neurontin).

Decision rationale: The MTUS considers Gabapentin as a first-line treatment for neuropathic pain and effective for the treatment of spinal cord injury, lumbar spinal stenosis, and post op pain. MTUS also recommends a trial of Gabapentin for complex regional pain syndrome. ODG states "Recommended Trial Period: One recommendation for an adequate trial with Gabapentin is three to eight weeks for titration, then one to two weeks at maximum tolerated dosage. (Dworkin, 2003) The patient should be asked at each visit as to whether there has been a change in pain or function. Current consensus based treatment algorithms for diabetic neuropathy suggests that if inadequate control of pain is found, a switch to another first-line drug is recommended." Additionally, ODG states that Gabapentin "has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain." Based on the clinical documentation provided, there is evidence of neuropathic type pain and radicular pain on exam and subjectively. The patient gets pain relief and functional improvement with the medication. As such, the request for Gralise ER: 600mg #120 is medically necessary.

Flexeril 10mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 64.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine, Medications for chronic pain, Antispasmodics Page(s): 41-42, 60-61, 64-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Cyclobenzaprine (Flexeril) and Other Medical Treatment Guidelines UpToDate, Flexeril.

Decision rationale: MTUS Chronic Pain Medical Treatment states for Cyclobenzaprine, "Recommended as an option, using a short course of therapy. The effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. (Browning, 2001) Treatment should be brief." The medical documents indicate that patient is far in excess of the initial treatment window and period. Additionally, MTUS outlines that "Relief of pain with the use of medications is generally temporary, and measures of the lasting benefit from this modality should include evaluating the effect of pain relief in relationship to improvements in function and increased activity. Before prescribing any medication for pain the following should occur: (1) determine the aim of use of the medication; (2) determine the potential benefits and adverse effects; (3) determine the patient's preference. Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) UpToDate "flexeril" also recommends, "Do not use longer than 2-3

weeks." Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine. ODG states regarding cyclobenzaprine, "Recommended as an option, using a short course of therapy. The addition of cyclobenzaprine to other agents is not recommended." As such, the request for Flexeril 10mg #30 is not medically necessary.