

Case Number:	CM15-0077941		
Date Assigned:	04/29/2015	Date of Injury:	05/04/2012
Decision Date:	05/26/2015	UR Denial Date:	04/15/2015
Priority:	Standard	Application Received:	04/23/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Physical Medicine & Rehabilitation

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 39 year old male, who sustained an industrial injury on 5/4/2012. He reported injury from a slip and fall. The injured worker was diagnosed as having lumbar radiculitis, lumbago and chronic pain. There is no record of a recent diagnostic study. Treatment to date has included medication management. In a progress note dated 4/6/2015, the injured worker complains of left leg pain. The treating physician is requesting Norco and Xanax.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325 mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page 79-80.

Decision rationale: Pain symptoms and clinical findings remain unchanged for this chronic injury. Submitted documents show no evidence that the treating physician is prescribing opioids

in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or returned to work status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury. In addition, submitted reports have not adequately demonstrated the specific indication to support for chronic opioid use without acute flare-up, new injuries, or progressive clinical deficits to support for chronic opioids outside recommendations of the guidelines. The Norco 10/325 mg #120 is not medically necessary and appropriate.

Xanax 1 mg #120: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines, page 24.

Decision rationale: There is no report of acute exacerbation or new injuries reported. Xanax Tablets (alprazolam) is indicated for the management of anxiety disorder. Anxiety or tension associated with the stress of everyday life usually does not require treatment with an anxiolytic. Alprazolam is an anti-anxiety medication in the benzodiazepine family which inhibits many of the activities of the brain as it is believed that excessive activity in the brain may lead to anxiety or other psychiatric disorders. Per the Chronic Pain Treatment Guidelines, benzodiazepines are not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks as chronic benzodiazepines are the treatment of choice in very few conditions and tolerance to hypnotic effects develops rapidly. The Xanax 1 mg #120 is not medically necessary and appropriate.