

Case Number:	CM15-0078127		
Date Assigned:	04/29/2015	Date of Injury:	11/11/2014
Decision Date:	05/26/2015	UR Denial Date:	04/16/2015
Priority:	Standard	Application Received:	04/23/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: Arizona, California
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 52 year old female who sustained an industrial injury on 11/11/14. The injured worker reported symptoms in the right shoulder. The injured worker was diagnosed as having a frozen shoulder, disorder of rotator cuff, full thickness rotator cuff tear, hip impingement syndrome, cervical strain, strain of left knee and strain of right wrist. Treatments to date have included acupuncture treatment, injections, and physical therapy. Currently, the injured worker complains of right shoulder discomfort. The plan of care was for a trigger point injection trial.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Trigger Point Injection Trial: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS ACOEM Page(s): 204. Decision based on Non-MTUS Citation ODG pain chapter and trigger point- pg 90.

Decision rationale: According to the guidelines, Invasive techniques have limited proven value. According to the ODG guidelines trigger point injections are not recommended in the absence of myofascial pain: Criteria for the use of Trigger point injections: Trigger point injections (TPI) with a local anesthetic with or without steroid may be recommended for the treatment of chronic low back or neck pain with myofascial pain syndrome (MPS) when all of the following criteria are met: (1) Documentation of circumscribed trigger points with evidence upon palpation of a twitch response as well as referred pain; (2) Symptoms have persisted for more than three months; (3) Medical management therapies such as ongoing stretching exercises, physical therapy, NSAIDs and muscle relaxants have failed to control pain; (4) Radiculopathy is not an indication (however, if a patient has MPS plus radiculopathy a TPI may be given to treat the MPS); (5) Not more than 3-4 injections per session; (6) No repeat injections unless a greater than 50% pain relief with reduced medication use is obtained for six weeks after an injection and there is documented evidence of functional improvement; (7) Frequency should not be at an interval less than two months; (8) Trigger point injections with any substance (e.g., saline or glucose) other than local anesthetic with or without steroid are not recommended; (9) There should be evidence of continued ongoing conservative treatment including home exercise and stretching. Use as a sole treatment is not recommended; (10) If pain persists after 2 to 3 injections the treatment plan should be re-examined as this may indicate an incorrect diagnosis, a lack of success with this procedure. In this case, the trigger point region was not well defined not spasms mentioned in the recent progress notes. Based on the above guidelines, a trigger point trial is not medically necessary.