

Case Number:	CM15-0078404		
Date Assigned:	04/29/2015	Date of Injury:	03/13/2012
Decision Date:	05/26/2015	UR Denial Date:	04/16/2015
Priority:	Standard	Application Received:	04/23/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The 45 year old male injured worker suffered an industrial injury on 03/13/2012. The diagnoses included lumbar disc herniation and lumbar strain. The diagnostics included lumbar magnetic resonance imaging. The injured worker had been treated with medications. On 7/18/2014 the treating provider reported the pain 7/10 with medications and 10/10 without medications radiating to the left leg from the low back with poor range of motion and acute muscle spasms. The treatment plan included MRI of the lumbosacral spine, Referral to neurosurgeon and Norco.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MRI of the lumbosacral spine: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304.

Decision rationale: The ACOEM chapter on low back complaints and special diagnostic studies states: Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computed tomography [CT] for bony structures). Relying solely on imaging studies to evaluate the source of low back and related symptoms carries a significant risk of diagnostic confusion (false positive test results) because of the possibility of identifying a finding that was present before symptoms began and therefore has no temporal association with the symptoms. Techniques vary in their abilities to define abnormalities (Table 12-7). Imaging studies should be reserved for cases in which surgery is considered or red-flag diagnoses are being evaluated. Because the overall false-positive rate is 30% for imaging studies in patients over age 30 who do not have symptoms, the risk of diagnostic confusion is great. There is no recorded presence of emerging red flags on the physical exam. There is evidence of nerve compromise on physical exam but there is not mention of consideration for surgery or complete failure of conservative therapy. For these reasons, criteria for imaging as defined above per the ACOEM have not been met. Therefore the request is not medically necessary.

Referral to neurosurgeon for consult and treatment: Overturned

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation ACOEM Practice Guidelines, Chapter 7: Independent Medical Examinations and Consultations, page 127.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 2 General Approach to Initial Assessment and Documentation, Chapter 3 Initial Approaches to Treatment.

Decision rationale: Per the ACOEM: The health practitioner may refer to other specialist if a diagnosis is uncertain or extremely complex, when psychosocial factors are present, or when the plan or course of care may benefit from additional expertise. A referral may be for 1. Consultation to aid in the diagnosis, prognosis, therapeutic management, determination of medical stability. The patient has ongoing back pain despite conservative therapy. The referral for a neurosurgeon would thus be medically necessary and approved.

Norco 10/325mg #180: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids
Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work; (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore criteria for the ongoing use of opioids have not been met and the request is not medically necessary.